

WEAKLY \mathcal{P} -SATURATED GRAPHS

MIECZYSLAW BOROWIECKI

AND

ELŻBIETA SIDOROWICZ

*Institute of Mathematics
University of Zielona Góra
65-246 Zielona Góra, Podgórna 50, Poland*

e-mail: M.Borowiecki@im.uz.zgora.pl

e-mail: E.Sidorowicz@im.uz.zgora.pl

Abstract

For a hereditary property \mathcal{P} let $k_{\mathcal{P}}(G)$ denote the number of forbidden subgraphs contained in G . A graph G is said to be *weakly \mathcal{P} -saturated*, if G has the property \mathcal{P} and there is a sequence of edges of \overline{G} , say e_1, e_2, \dots, e_l , such that the chain of graphs $G = G_0 \subset G_0 + e_1 \subset G_1 + e_2 \subset \dots \subset G_{l-1} + e_l = G_l = K_n$ ($G_{i+1} = G_i + e_{i+1}$) has the following property: $k_{\mathcal{P}}(G_{i+1}) > k_{\mathcal{P}}(G_i)$, $0 \leq i \leq l-1$.

In this paper we shall investigate some properties of weakly saturated graphs. We will find upper bound for the minimum number of edges of weakly \mathcal{D}_k -saturated graphs of order n . We shall determine the number $\text{wsat}(n, \mathcal{P})$ for some hereditary properties.

Keywords: graph, extremal problems, hereditary property, weakly saturated graphs.

1991 Mathematics Subject Classification: 05C35.

1. INTRODUCTION AND NOTATION

We consider finite undirected graphs without loops or multiple edges. A graph G has a vertex set $V(G)$ and an edge set $E(G)$. Let $v(G)$, $e(G)$ denote the number of vertices and the number of edges of G , respectively. We say that G *contains* H whenever G contains a subgraph isomorphic to H .

The degree of $v \in V(G)$ is denoted by $d_G(v)$. The number of edges of a path is called the *length* of the path.

Let \mathcal{I} denote the class of all graphs with isomorphic graphs being regarded as equal. If \mathcal{P} is a proper nonempty subclass of \mathcal{I} , then \mathcal{P} will also denote the property of being in \mathcal{P} . We shall use the terms *class of graphs* and *property of graphs* interchangeably.

A property \mathcal{P} is called *hereditary* if every subgraph of a graph G with property \mathcal{P} also has property \mathcal{P} .

We list some properties to introduce the necessary notation which will be used in the paper. Let k be a non-negative integer.

$$\mathcal{O} = \{G \in \mathcal{I} : G \text{ is totally disconnected}\},$$

$$\mathcal{O}_k = \{G \in \mathcal{I} : \text{each component of } G \text{ has at most } k+1 \text{ vertices}\},$$

$$\mathcal{I}_k = \{G \in \mathcal{I} : G \text{ contains no subgraph isomorphic to } K_{k+2}\},$$

$$\mathcal{S}_k = \{G \in \mathcal{I} : \Delta(G) \leq k\},$$

$$\mathcal{D}_k = \{G \in \mathcal{I} : G \text{ is } k\text{-degenerated, i.e., } \delta(H) \leq k \text{ for any } H \leq G\},$$

$$\mathcal{W}_k = \{G \in \mathcal{I} : \text{the length of the longest path in } G \text{ is at most } k\}.$$

Let \mathcal{P} be a nontrivial hereditary property. Then there is a nonnegative integer $c(\mathcal{P})$, called the *completeness* of \mathcal{P} , such that $K_{c(p)+1} \in \mathcal{P}$ but $K_{c(p)+2} \notin \mathcal{P}$. Obviously

$$c(\mathcal{O}_k) = c(\mathcal{I}_k) = c(\mathcal{S}_k) = c(\mathcal{D}_k) = c(\mathcal{W}_k) = k.$$

For a hereditary property \mathcal{P} the set of all *minimal forbidden subgraphs* of \mathcal{P} is defined by

$$F(\mathcal{P}) = \{G \in \mathcal{I} : G \notin \mathcal{P} \text{ but each proper subgraph } H \text{ of } G \text{ belongs to } \mathcal{P}\}.$$

A graph is called *\mathcal{P} -maximal* if it does not contain any forbidden subgraph but it will contain a forbidden subgraph when any new edge is added to the graph. Let $M(\mathcal{P})$ be the set of all \mathcal{P} -maximal graphs. The set of \mathcal{P} -maximal graphs of order n is denoted by $M(n, \mathcal{P})$.

Many problems of extremal graph theory can be formulated as follows: What is the maximum (minimum) number of edges in a \mathcal{P} -maximal graph

of order n ? For a given hereditary property \mathcal{P} we define those two numbers in the following manner:

$$\begin{aligned} \text{ex}(n, \mathcal{P}) &= \max\{e(G) : G \in \text{M}(n, \mathcal{P})\}, \\ \text{sat}(n, \mathcal{P}) &= \min\{e(G) : G \in \text{M}(n, \mathcal{P})\}. \end{aligned}$$

The set of all \mathcal{P} -maximal graphs of order n with exactly $\text{ex}(n, \mathcal{P})$ edges is denoted by $\text{Ex}(n, \mathcal{P})$. The members of $\text{Ex}(n, \mathcal{P})$ are called \mathcal{P} -*extremal* graphs. By the symbol $\text{Sat}(n, \mathcal{P})$ is denoted the set of all \mathcal{P} -maximal graphs of order n with $\text{sat}(n, \mathcal{P})$ edges. These graphs are called \mathcal{P} -*saturated*.

The most famous Turán's Theorem [6] establishes the number of edges of \mathcal{I}_k -extremal graphs. On the other hand, Erdős, Hajnal and Moon [2] calculated the number $\text{sat}(n, \mathcal{I}_k)$.

Bollobás [1] introduced the concept of a weakly k -saturated graph. Consider a graph of order n and add all those edges which are the only missing edge of complete graph of order k (i.e., we add the edge e if there are k such vertices of the graph, that the graph contains all the edges spanned by these k vertices, saving e). If by repeating this process a sufficient number of times the complete graph of order n is obtained, the original graph will be called *weakly k -saturated*.

Bollobás showed that if a graph G of order n is weakly k -saturated (for $3 \leq k \leq 7$) with the minimum number of edges then $e(G) = (k-2)n - \binom{k-1}{2}$. In the general case (i.e., for $k \geq 3$) the equality has been proved by Kalai [5].

Let \mathcal{P} be a hereditary property and let $k_{\mathcal{P}}(G)$ denote the number of forbidden subgraphs contained in G . A graph G is said to be *weakly \mathcal{P} -saturated*, if G has the property \mathcal{P} and there is a sequence of edges of \overline{G} , say e_1, e_2, \dots, e_l , such that the chain of graphs $G = G_0 \subset G_0 + e_1 \subset G_1 + e_2 \subset \dots \subset G_{l-1} + e_l = G_l = K_n$ ($G_{i+1} = G_i + e_{i+1}$) has the following property: $k_{\mathcal{P}}(G_{i+1}) > k_{\mathcal{P}}(G_i)$, $0 \leq i \leq l-1$. This sequence of edges will be called the *complementary sequence of G with respect to \mathcal{P}* or briefly the *complementary sequence* if it does not lead us to misunderstanding.

According to our terminology a weakly k -saturated graph is called weakly \mathcal{I}_{k-2} -saturated.

Let us denote a set of all weakly \mathcal{P} -saturated graphs of order n by $\text{WSat}(n, \mathcal{P})$. Let the minimum and the maximum number of edges in a

graph of $\text{WSat}(n, \mathcal{P})$ be denoted by

$$\text{wsat}(n, \mathcal{P}) = \min\{e(G) : G \in \text{WSat}(n, \mathcal{P})\},$$

$$\text{wex}(n, \mathcal{P}) = \max\{e(G) : G \in \text{WSat}(n, \mathcal{P})\}.$$

From Theorem of Kalai and Theorem of Erdős, Hajnal, Moon it follows that $\text{wsat}(n, \mathcal{I}_k) = \text{sat}(n, \mathcal{I}_k)$. In Section 2 we shall describe a hereditary property \mathcal{P} such that $\text{wsat}(n, \mathcal{P}) < \text{sat}(n, \mathcal{P})$. We will also investigate some properties of weakly saturated graphs. In Section 3 examples of weakly \mathcal{D}_k -saturated graphs and an upper bound for the number $\text{wsat}(n, \mathcal{D}_k)$ will be given. In Section 4 we shall determine the number $\text{wsat}(n, \mathcal{P})$ for some hereditary properties.

2. SOME PROPERTIES OF WEAKLY \mathcal{P} -SATURATED GRAPHS

From the definition of weakly \mathcal{P} -saturated graphs it follows that any \mathcal{P} -maximal graph is weakly \mathcal{P} -saturated. First we prove that the maximum number of edges of weakly \mathcal{P} -saturated graphs is equal to the maximum number of edges of \mathcal{P} -maximal graphs.

Theorem 1. *Let $n \geq 1$. If \mathcal{P} is a hereditary property, then $\text{wex}(n, \mathcal{P}) = \text{ex}(n, \mathcal{P})$.*

Proof. Every \mathcal{P} -maximal graph is weakly \mathcal{P} -saturated. Thus $\text{wex}(n, \mathcal{P}) \geq \text{ex}(n, \mathcal{P})$. On the other hand, if a graph of order n has more than $\text{ex}(n, \mathcal{P})$ edges then it contains a forbidden subgraph. Hence $\text{wex}(n, \mathcal{P}) \leq \text{ex}(n, \mathcal{P})$. ■

Any non-negative integer valued function $f : \mathcal{I} \rightarrow N$ is called the *graph invariant* (*invariant*, for short). For a hereditary property \mathcal{P} let us define the number

$$f(\mathcal{P}) = \min\{f(H) : H \in \mathcal{F}(\mathcal{P})\}.$$

Theorem 2. *Let $f(G)$ be an invariant satisfying:*

- (1) $f(H) \leq f(G)$ for $H \subseteq G$,
- (2) $f(G + e) \leq f(G) + 1$ for $e \in E(\overline{G})$.

Then for any graph $G \in \text{WSat}(n, \mathcal{P})$ with $n \geq c(\mathcal{P}) + 2$, we have

$$f(G) \geq f(\mathcal{P}) - 1.$$

Proof. From the definition of weakly \mathcal{P} -saturated graphs, it follows that there is an edge $e \in E(\overline{G})$ and a graph $F \in \mathcal{F}(\mathcal{P})$ such that $F \subseteq G + e$. Thus $f(\mathcal{P}) \leq f(F) \leq f(G + e) \leq f(G) + 1$. ■

The chromatic number and the clique number are examples of invariant satisfying assumptions of Theorem 2. The edge connectivity $\lambda(G)$ does not satisfy the assumption (1) of Theorem 2, but we shall prove that for $G \in \text{WSat}(n, \mathcal{P})$ the inequality $\lambda(G) \geq \lambda(\mathcal{P}) - 1$ also holds.

Theorem 3. *Let $\lambda(\mathcal{P}) = \lambda > 0$ and $G \in \text{WSat}(n, \mathcal{P})$. Then*

$$\lambda(G) \geq \lambda - 1.$$

Proof. Let S be an edge cutset of G such that $\lambda(G) = |S|$. Let G' , G'' be two components of $G - S$. Since G is weakly \mathcal{P} -saturated, it follows that there is a complementary sequence e_1, e_2, \dots, e_l of G . Let e_i be the first edge of the sequence e_1, e_2, \dots, e_l , which joins a vertex of G' with a vertex of G'' . Let F denote a subgraph of $G_{i-1} + e_i$, which contains the edge e_i and is isomorphic with some graph of $\mathcal{F}(\mathcal{P})$. Then the set $S \cup \{e_i\}$ is an edge cutset of F . Thus $\lambda \leq \lambda(F) \leq |S| + 1 = \lambda(G) + 1$. ■

From the next theorem it follows that the behaviour of $\text{wsat}(n, \mathcal{P})$ is not monotone in general.

Theorem 4. *Let \mathcal{P} be the hereditary property such that $\mathcal{F}(\mathcal{P}) = \{2K_2\}$. Then*

$$\text{wsat}(n, \mathcal{P}) = \begin{cases} 3, & \text{for } n = 4, \\ 1, & \text{for } n \geq 5. \end{cases}$$

Proof. It is easy to see that there is no weakly \mathcal{P} -saturated graph of order 4 with two edges. Since the graphs $K_{1,3}$ and $K_3 \cup K_1$ are weakly \mathcal{P} -saturated, we have $\text{wsat}(4, \mathcal{P}) = 3$.

If $n \geq 5$ then $K_2 \cup (n-2)K_1$ is a weakly \mathcal{P} -saturated graph. By adding (as long as possible) an edge joining two vertices of $(n-2)K_1$ we obtain two independent edges, i.e., $2K_2$, and results in K_{n-2} . Since $n-2 \geq 3$, it follows that every vertex of K_2 (in the original graph), we can join with every vertex of just obtained K_{n-2} . ■

From Theorem of Kalai and Theorem of Erdős, Hajnal and Moon, it follows that $\text{wsat}(n, \mathcal{I}_k) = \text{sat}(n, \mathcal{I}_k)$. Such equality also holds for the property \mathcal{D}_1 .

Theorem 5. *Let $n \geq 1$. Then*

$$\text{sat}(n, \mathcal{D}_1) = \text{wsat}(n, \mathcal{D}_1) = n - 1.$$

Proof. Since $F(\mathcal{D}_1) = \{C_p : p \geq 3\}$, $\lambda(\mathcal{D}_1) = 2$ and every tree is weakly \mathcal{D}_1 -saturated, it follows that $\text{wsat}(n, \mathcal{D}_1) \leq n - 1$. From Theorem 3 we have $\lambda(G) \geq 1$ for $G \in \text{WSat}(n, \mathcal{D}_1)$ then $\text{wsat}(n, \mathcal{D}_1) \geq n - 1$. Thus $\text{wsat}(n, \mathcal{D}_1) = n - 1$. Since the only \mathcal{D}_1 -maximal graphs are trees, we have $\text{sat}(n, \mathcal{D}_1) = n - 1$. ■

The next theorem describes a hereditary property \mathcal{P} for which the minimum number of edges of weakly \mathcal{P} -saturated graphs of order n is less than the number of edges of \mathcal{P} -saturated graphs of order n .

Theorem 6. *Let \mathcal{P} be the hereditary property such that $\text{ex}(n, \mathcal{P}) = \text{sat}(n, \mathcal{P})$, $\lambda(\mathcal{P}) = \lambda(H_0) = 1$, $H_0 \in F(\mathcal{P})$ and every \mathcal{P} -maximal graph is connected. Then $\text{wsat}(n, \mathcal{P}) < \text{sat}(n, \mathcal{P})$, $n \geq v(H_0)$.*

Proof. Let $H_0 \in F(\mathcal{P})$ with $\lambda(H_0) = 1$ and let e be a cutedge of H_0 . Denote by H_1, H_2 components of $H_0 - e$. Let $v(H_1) = n_1$, $v(H_2) = n_2$. We define the graph $G = G_1 \cup G_2$ of order n assuming that $v(G_1) = n_1$, $v(G_2) = n - n_1$ and for $i = 1, 2$, G_i is \mathcal{P} -maximal. Obviously $n - n_1 \geq n_2$. Since all forbidden subgraphs are connected it follows that the graph G has property \mathcal{P} . Defined graph G is not connected, then by the assumption of the theorem, G is not \mathcal{P} -maximal. Thus $e(G) < \text{ex}(n, \mathcal{P}) = \text{sat}(n, \mathcal{P})$.

On the other hand, we will show that the graph G is weakly \mathcal{P} -saturated. Since each component of G is a \mathcal{P} -maximal graph, it follows that if we add any edge of \overline{G} which joins two vertices of the same component we obtain a new forbidden subgraph containing the edge e . After adding all missing edges of each component we obtain the graph being a sum of complete graphs. Then each edge, which joins a vertex of the component of order n_1 with a vertex of the component of order $n - n_1$, belongs to a subgraph isomorphic to H_0 . Thus the graph G is weakly \mathcal{P} -saturated and $e(G) \geq \text{wsat}(n, \mathcal{P})$. Hence $\text{wsat}(n, \mathcal{P}) < \text{sat}(n, \mathcal{P})$. ■

In the next section we will show that the assumptions of Theorem 6 for the property \mathcal{D}_k ($k \geq 2$) holds.

3. WEAKLY \mathcal{D}_k -SATURATED GRAPHS

The set of minimal forbidden subgraphs for property \mathcal{D}_k was characterized by Mihók [4]. To describe the set $F(\mathcal{D}_k)$ we need some more notations. For a nonnegative integer k and a graph G , we denote the set of all vertices of G of degree $k + 1$ by $M(G)$. If $S \subseteq V(G)$ is a cutset of vertices of G and G_1, \dots, G_s , $s \geq 2$ are the components of $G - S$, then the graph $G - V(G_i)$ is denoted by H_i , $i = 1, \dots, s$.

Theorem 7. [4] *A graph G belongs to $F(\mathcal{D}_k)$ if and only if G is connected, $\delta(G) \geq k + 1$, $V(G) - M(G)$ is an independent set of vertices of G and for each cutset $S \subset V(G) - M(G)$ we have that $\delta(H_i) \leq k$ for each $i = 1, \dots, s$.*

Let us present some useful examples of $F(\mathcal{D}_k)$.

Example 1. Let H_k , $k \geq 2$, be the graph such that $V(H_k) = \{x_1, \dots, x_k, y_1, \dots, y_k, v_1, v_2, w_1, w_2\}$ with the following properties: vertices x_1, \dots, x_k and y_1, \dots, y_k induce two complete graphs and $v_i w_i, v_i x_j, w_i y_j \in E(H_k)$ for $i = 1, 2$, $j = 1, \dots, k$.





Figure 3.1. The graph H_k for $k = 2$

Example 2. Let H'_k , $k \geq 2$, be the graph such that $V(H'_k) = \{x_1, \dots, x_k, y_1, \dots, y_k, v_1, v_2, v_3, w_1, w_2, w_3\}$ with the following properties: vertices x_1, \dots, x_k and y_1, \dots, y_k induce two graphs obtained from K_k by removing $\lfloor \frac{k}{2} \rfloor$ independent edges and $v_i x_j, w_i y_j \in E(H'_k)$ for $i = 1, 2, 3$, $j = 1, \dots, k$, and $v_1 w_1, v_2 v_3, w_2 w_3 \in E(H'_k)$.

Figure 3.2. The graph H'_k for $k = 2$

By Example 2 we have that $\lambda(\mathcal{D}_k) = 1$ for $k \geq 2$. Since \mathcal{D}_k -maximal graphs are connected and $\text{sat}(n, \mathcal{D}_k) = \text{ex}(n, \mathcal{D}_k)$ (see e.g. [3]), it follows that the assumptions of Theorem 6 holds. Then we immediately have

Corollary 8. $\text{wsat}(n, \mathcal{D}_k) < \text{sat}(n, \mathcal{D}_k)$ for $n \geq 2(k+3)$, $k \geq 2$.

To determine upper bound for the number $\text{wsat}(n, \mathcal{D}_k)$ we need the following lemma.

Lemma 9. *Let $k \geq 2$. Then the graph $H_k - v_2w_2$ is weakly \mathcal{D}_k -saturated.*

Proof. Put $G = H_k - v_2w_2$. If the edge v_2w_2 is added to G then $G = H_k \in \mathcal{F}(\mathcal{D}_k)$ is obtained. If we add v_1v_2 or w_1w_2 to H_k then we obtain the graph K_{k+2} which belongs to $\mathcal{F}(\mathcal{D}_k)$. After adding the edge x_iy_j , $(1 \leq i, j \leq k)$, edges $(E(G) \cup \{v_1v_2, w_1w_2, x_iy_j\}) - \{v_1x_i, w_1y_j\}$ induce H_k . Now we can add the edge v_1y_j , $1 \leq j \leq k$ since edges $(E(G) \cup \{v_2w_2, w_1w_2, v_1y_j\}) - \{w_2y_j, v_1w_1\}$ induce H_k . If we add the edge v_2w_j ($1 \leq j \leq k$), we obtain the graph H_k induced by $(E(G) \cup \{w_1w_2, v_2w_j\}) - \{w_1y_j\}$. In a similar manner we can show that if we add edges x_iw_1 and x_iw_2 ($1 \leq i \leq k$), a new forbidden subgraph appears. The last two edges v_1w_2 , v_2w_1 we can add because edges $(E(G) \cup \{x_1y_1, v_1w_2, v_1v_2, w_1w_2\}) - \{x_1v_1, w_2y_1, v_1w_1\}$ and $(E(G) \cup \{x_1y_1, v_2w_1, v_1v_2, w_1w_2\}) - \{x_1v_2, w_1y_1, v_1w_1\}$ induce H_k . ■

Theorem 10. *Let $k \geq 2$ and $n = 2(k+2)q+r$, where $q \geq 1$, $0 \leq r \leq 2k+3$. Then*

$$\text{wsat}(n, \mathcal{D}_k) \leq \begin{cases} \frac{(k+2)(k+1)-1}{2(k+2)}n, & \text{for } r = 0, \\ \frac{(k+2)(k+1)-1}{2(k+2)}(n - r - (k+2)) + \\ \quad (r + k + 2)k - \binom{k+1}{2}, & \text{for } 0 < r < k+3, \\ \frac{(k+2)(k+1)-1}{2(k+2)}(n - r) + rk - \binom{k+1}{2}, & \text{for } r \geq k+3. \end{cases}$$

Proof. To prove the theorem it is enough to show that there is a weakly \mathcal{D}_k -saturated graph G of order n with such number of edges. Let $k \geq 2$ and $n = 2(k+2)q + r$, where $q \geq 1$, $0 \leq r \leq 2k+3$. Put $G' = H_k - v_2w_2$. If $r \geq k+3$, then $G = qG' \cup H$, where $H \in \mathcal{M}(r, \mathcal{D}_k)$. If $0 \leq r < k+3$, then $G = (q-1)G' \cup H$, where $H \in \mathcal{M}(2(k+2)+r, \mathcal{D}_k)$. If $r = 0$, then $G = qG'$. By Lemma 9 it follows that each component of G is a weakly \mathcal{D}_k -saturated graph. Then we can add edges in each component of G to obtain a complete graph. After having added those edges we can join any vertices of two different components. ■

4. THE NUMBER $\text{wsat}(n, \mathcal{P})$ FOR SOME HEREDITARY PROPERTIES

In this section we will calculate the minimum number of edges of weakly saturated graphs for some hereditary properties.

Theorem 11. *Let $k \geq 1$ and $n \geq k+2$. Then*

$$\text{WSat}(n, \mathcal{O}_k) \supseteq \{T_r \cup T_s \cup tT_1 : r+s = k+2, r+s+t = n \text{ and } T_i \text{ is an arbitrary tree of order } i\}$$

and

$$\text{wsat}(n, \mathcal{O}_k) = k.$$

Proof. First we prove that the graph $G = T_r \cup T_s \cup tT_1$, where $r+s = k+2$, $r+s+t = n$ is weakly \mathcal{O}_k -saturated. If we add an edge of \overline{G} , which joins a vertex of T_r and a vertex of T_s then we obtain a tree of order $k+2$, i.e., we obtain a forbidden subgraph for property \mathcal{O}_k . If we join a vertex of the subgraph tT_1 with a vertex of the obtained tree of order $k+2$ we have a connected graph of order $k+3$. Thus new edge belongs to a tree of order $k+2$. Repeating this process we obtain a connected graph of order n in which each vertex of tT_1 is adjacent with any vertex of the tree of order $k+2$. Since for each edge of the complement of a connected graph there is a spanning tree which contains this edge, it follows that G is weakly \mathcal{O}_k -saturated. Hence $\text{wsat}(n, \mathcal{O}_k) \leq e(G) = k$.

On the other hand, let G be a graph such that $G \in \text{WSat}(n, \mathcal{O}_k)$ and $e(G) = \text{wsat}(n, \mathcal{O}_k)$. Let e_1 be the first edge such that $G + e_1$ contains a forbidden subgraph, i.e., the graph $G + e_1$ contains a tree of order $k+2$. Thus $\text{wsat}(n, \mathcal{O}_k) = e(G) \geq k$. ■

The proof of the next theorem is very similar to the proof of Theorem 11, then it is omitted.

Theorem 12. *Let $k \geq 1$ and $n \geq k + 2$. Then*

$$\text{WSat}(n, \mathcal{W}_k) \supseteq \{P_r \cup P_s \cup tP_1 : r + s = k + 2, r + s + t = n\}$$

and

$$\text{wsat}(n, \mathcal{W}_k) = k.$$

It is easy to see that the graphs $K_{k+1} + tK_1$, where $k + 1 + t = n$ are weakly \mathcal{S}_k -saturated. There are some other weakly \mathcal{S}_k -saturated graphs of order n . For example the graph G_1 (Figure 4.1) is weakly \mathcal{S}_2 -saturated and the graph G_2 (Figure 4.1) is weakly \mathcal{S}_3 -saturated.




Figure 4.1. The graphs G_1 and G_2

Theorem 13. *Let $n \geq k + 2 \geq 4$. Then*

$$\text{wsat}(n, \mathcal{S}_k) = \binom{k+1}{2}.$$

Proof. Let G be a weakly \mathcal{S}_k -saturated graph of order n with the minimum number of edges. Then there is a complementary sequence e_1, e_2, \dots, e_l of G . Let $e_1 = u_1v_1$ and $d_G(u_1) = k$. Let $e_{f(1)}, \dots, e_{f(t_1)}$ be the subsequence of e_1, e_2, \dots, e_l such that every edge $e_{f(i)}$, ($1 \leq i \leq t_1$) is adjacent with the vertex u_1 . If in the graph $G' = ((G + e_{f(1)}) + e_{f(2)}) + \dots + e_{f(t_1)}$ there is no vertex of degree less than k then let $e_{f(1)}, e_{f(2)}, \dots, e_{f(l)}$ be the new sequence of edges of $E(\overline{G})$ with the following property: $e_{f(1)}, \dots, e_{f(t_1)}$ is the subsequence of e_1, e_2, \dots, e_l such that every edge $e_{f(i)}$, ($1 \leq i \leq t_1$) is adjacent with the vertex u_1 and $e_{f(t_1)+1}, \dots, e_{f(l)}$ is the subsequence of e_1, e_2, \dots, e_l such that any edge $e_{f(i)}$, ($t_1 \leq i \leq l$) is not adjacent with the vertex u_1 . If in the graph G' there is a vertex of degree less than k

then let $e_{f(t_1+1)}$ be the first edge of e_1, e_2, \dots, e_l , which is not adjacent with the vertex u_1 . Let $e_{f(t_1+1)} = u_2 v_2$ and u_2 be a vertex of G' such that $d_{G'}(u_2) \geq k$ and $u_1 \neq u_2$. Let $e_{f(t_1+1)}, \dots, e_{f(t_2)}$ denote edges of $\{e_1, e_2, \dots, e_l\} - \{e_{f(1)}, \dots, e_{f(t_1)}, e_{f(t_1)+1}\}$ which are adjacent with the vertex u_2 . If in the graph $G'' = ((G' + e_{f(t_1)+1}) + e_{f(t_1)+2}) + \dots + e_{f(t_2)}$ there is no vertex of degree less than k we form a new sequence of edges of $E(\overline{G})$, $e_{f(1)}, e_{f(2)}, \dots, e_{f(l)}$ with the following property: $e_{f(1)}, \dots, e_{f(t_1)}$ is a subsequence of e_1, e_2, \dots, e_l such that every edge $e_{f(i)}$, $(1 \leq i \leq t_1)$ is adjacent with the vertex u_1 and $e_{f(t_1)+1}, \dots, e_{f(t_2)}$ is a subsequence of e_1, e_2, \dots, e_l such that every edge $e_{f(i)}$, $(t_1 < i \leq t_2)$ is adjacent with the vertex u_2 and $e_{f(t_2)+1}, \dots, e_{f(l)}$ is the subsequence of e_1, e_2, \dots, e_l such that any edge $e_{f(i)}$, $(t_2 < i \leq l)$ is not adjacent with the vertex u_1 and u_2 . If in the graph G'' there is a vertex of degree less than k , we will repeat this steps until we will obtain a new sequence $e_{f(1)}, e_{f(2)}, \dots, e_{f(l)}$ of edges of \overline{G} . With this sequence of edges $e_{f(1)}, e_{f(2)}, \dots, e_{f(l)}$ is related a sequence of vertices u_1, u_2, \dots, u_r . It is easy to see that $r \leq k$, because after k steps there is no vertex of degree less than k . Then for the vertex $u_t \in \{u_1, \dots, u_r\}$ we have

$$(1) \quad d_G(u_t) + t - 1 - |N_G(u_t) \cap \{u_1, \dots, u_{t-1}\}| \geq k,$$

for the vertex $x \in V(G) - \{u_1, \dots, u_r\}$ we have

$$(2) \quad d_G(x) + r - |N_G(x) \cap \{u_1, \dots, u_r\}| \geq k.$$

Thus

$$\begin{aligned} e(G) &\geq \sum_{1 \leq t \leq r} (d_G(u_t) - |N_G(u_t) \cap \{u_1, \dots, u_{t-1}\}|) \\ &\quad + \frac{1}{2} \sum_{x \in V(G) - \{u_1, \dots, u_r\}} (d_G(x) - |N_G(x) \cap \{u_1, \dots, u_r\}|) \\ &\geq \sum_{1 \leq t \leq r} (k + 1 - t) + \frac{1}{2} (n - r)(k - r). \end{aligned}$$

The right side of inequality achieves the minimum for $r = k$. Thus

$$e(G) \geq \sum_{1 \leq t \leq r} (k + 1 - t) = \frac{1}{2} (k + 1)k.$$

On the other hand, the graph $K_{k+1} \cup (n - k - 1)K_1$ is weakly \mathcal{S}_k -saturated. Thus $\text{wsat}(n, \mathcal{S}_k) \leq \binom{k+1}{2}$. \blacksquare

In the next theorem we determine the number $\text{wsat}(n, \mathcal{P})$ for a hereditary property with one forbidden subgraph which is a cycle of odd length.

Theorem 14. *Let $k \geq 1$ and $n \geq 2k + 2$. If \mathcal{P} is the hereditary property such that $\text{F}(\mathcal{P}) = \{C_{2k+1}\}$, then $\text{wsat}(n, \mathcal{P}) = n - 1$.*

Proof. Since $\lambda(\mathcal{P}) = 2$, by Theorem 3 it follows that every weakly \mathcal{P} -saturated graph is connected. Then $\text{wsat}(n, \mathcal{P}) \geq n - 1$. To prove that the inequality $\text{wsat}(n, \mathcal{P}) \leq n - 1$ holds it is sufficient to show that there is a weakly \mathcal{P} -saturated graph of order n with $n - 1$ edges.

Let us show first that P_{2k+2} is a weakly \mathcal{P} -saturated graph. Let $V(P_{2k+2}) = \{v_1, \dots, v_{2k+2}\}$ and $d(v_1) = d(v_{2k+2}) = 1$. It is easy to see that if we add the edge $v_1 v_{2k+1}$ then we obtain a cycle of order $2k + 1$. Similarly if we add the edge $v_2 v_{2k+2}$ a new cycle of order $2k + 1$ appears. Now we can add the edge $v_1 v_4$. The edge $v_1 v_4$ belongs to the cycle $v_1, v_2, v_{2k+2}, v_{2k+1}, \dots, v_4, v_1$. To prove that if we add any edge $v_1 v_{2t}$ then a new cycles of order $2k + 1$ appears we will use induction on t . This is true for $t = 1, 2$. When the edges $v_1 v_{2i}$ for $i < t$ are added the vertices $v_1, v_{2t-2}, v_{2t-3}, \dots, v_2, v_{2k+2}, v_{2k+1}, \dots, v_{2t}, v_1$ induce a cycle of order $2k + 1$ which contains the edge $v_1 v_{2t}$. In the same manner, after having added edges $v_1 v_{2i+1}$ for $k \geq i > t$ we can add the edge $v_1 v_{2t+1}$. A new cycle $v_1, v_{2t+3}, \dots, v_{2k+2}, v_2, v_3, \dots, v_{2t+1}, v_1$ of order $2k + 1$ appears. Finally the vertex v_1 with all vertices of P_{2k+2} is joined. Similarly we can join each vertex v_t ($2 \leq t \leq 2k + 2$) with all vertices of P_{2k+2} . Thus we obtain a graph K_{2k+2} . Hence P_{2k+2} is a weakly \mathcal{P} -saturated graph.

Let G be the graph of order $n \geq 2k + 2$ with the following properties: G contains an induced path of order $2k + 2$, the remaining vertices of G form an independent set and each vertex of this set is adjacent with exactly one vertex of the path. Since the path of order $2k + 2$ is weakly \mathcal{P} -saturated, it follows that the graph G is weakly \mathcal{P} -saturated. Hence $\text{wsat}(n, \mathcal{P}) \leq n - 1$. ■

In order to determine the number $\text{wsat}(n, \mathcal{P})$ for hereditary property such that $F(\mathcal{P}) = \{C_{2k}\}$ we need the following lemma.

Lemma 15. *Let $k \geq 2$ and \mathcal{P} be the hereditary property such that $F(\mathcal{P}) = \{C_{2k}\}$, and G be a bipartite graph of order $n \geq 2k + 1$. Then $G \notin \text{WSat}(n, \mathcal{P})$.*

Proof. On the contrary, suppose that there is a weakly \mathcal{P} -saturated bipartite graph G of order n . Let e_1, e_2, \dots, e_l be a complementary sequence of G . Let $e_i = xy$ be the first edge of the sequence e_1, e_2, \dots, e_l such that its ends x, y belong to the same colour class of G . (Notice, that the colour classes of G are uniquely determined because of connectivity of G .) Since the edge e_i belongs to an even cycle C_{2k} then there is an edge e_j , $j < i$ of this cycle (and the sequence given above) with both ends in one colour class which is impossible. ■

Theorem 16. *Let $k \geq 2$ and $n \geq 2k + 1$. Let \mathcal{P} be the hereditary property such that $F(\mathcal{P}) = \{C_{2k}\}$. Then*

$$\text{wsat}(n, \mathcal{P}) = n.$$

Proof. Let $G \in \text{WSat}(n, \mathcal{P})$. By Theorem 3 and Lemma 15 it follows that G is connected and contains an odd cycle. Thus $\text{wsat}(n, \mathcal{P}) \geq n$.

To prove that the inequality $\text{wsat}(n, \mathcal{P}) \leq n$ holds it is sufficient to show that there is a weakly \mathcal{P} -saturated graph of order n with n edges. First we prove that C_{2k+1} is a weakly \mathcal{P} -saturated graph. Let $V(C_{2k+1}) = \{v_1, v_2, \dots, v_{2k+1}\}$. It is easy to see that if we add the edge v_1v_3 or the edge v_2v_{2k+1} , a cycle (containing this edge) of order $2k$ appears. To prove that if we add any edge v_1v_t ($3 \leq t \leq 2k$) then we obtain a new cycle of order $2k$ we use induction on t . This is true for $t = 3$. After adding edges v_1v_i for $3 \leq i < t$ the vertices $v_1, v_{t-2}, v_{t-3}, \dots, v_2, v_{2k+1}, v_{2k}, \dots, v_t, v_1$ induce a cycle of order $2k$ which contains the edge v_1v_t . Then the vertex v_1 can be joined with all vertices of C_{2k+1} . In the similar manner we can show that we can join any vertex $v_t \in V(C_{2k+1})$ with all vertices of C_{2k+1} . Hence C_{2k+1} is weakly \mathcal{P} -saturated.

Let G be the graph with the following properties: G contains an induced cycle of order $2k + 1$, remaining vertices of G form an independent set and each vertex of this set is adjacent with exactly one vertex of the cycle. Since the cycle of order $2k + 1$ is weakly \mathcal{P} -saturated (can be extended to K_{2k+1}), it follows that the graph G also has this property, i.e., G is weakly \mathcal{P} -saturated. Hence $\text{wsat}(n, \mathcal{P}) \leq n$. ■

REFERENCES

- [1] B. Bollobás, *Weakly k -saturated graphs*, in: H. Sachs, H.-J. Voss and H. Walther, eds, Proc. Beiträge zur Graphentheorie, Manebach, 9–12 May, 1967 (Teubner Verlag, Leipzig, 1968) 25–31.
- [2] P. Erdős, A. Hajnal and J.W. Moon, *A Problem in Graph Theory*, Amer. Math. Monthly **71** (1964) 1107–1110.
- [3] R. Lick and A. T. White, *k -degenerated graphs*, Canadian J. Math. **22** (1970) 1082–1096.
- [4] P. Mihók, *On graphs critical with respect to vertex partition numbers*, Discrete Math. **37** (1981) 123–126.

- [5] G. Kalai, *Weakly saturated graphs are rigid*, Annals of Discrete Math. **20** (1984) 189–190.
- [6] P. Turán, *On the Theory of Graphs*, Colloq. Math. **3** (1954) 19–30.

Received 20 August 2000

Revised 3 December 2001