

## ON GRAPHS $G$ FOR WHICH BOTH $G$ AND $\overline{G}$ ARE CLAW-FREE

SHINYA FUJITA

*Department of Mathematics*

*Keio University*

*Hiyoshi, Kohoku-ku, Yokohama, 223-8522, Japan*

**e-mail:** shinyaa@comb.math.keio.ac.jp

### Abstract

Let  $G$  be a graph with  $|V(G)| \geq 10$ . We prove that if both  $G$  and  $\overline{G}$  are claw-free, then  $\min\{\Delta(G), \Delta(\overline{G})\} \leq 2$ . As a generalization of this result in the case where  $|V(G)|$  is sufficiently large, we also prove that if both  $G$  and  $\overline{G}$  are  $K_{1,t}$ -free, then  $\min\{\Delta(G), \Delta(\overline{G})\} \leq r(t-1, t) - 1$  where  $r(t-1, t)$  is the Ramsey number.

**Keywords:** claw-free, complement, maximum degree.

**2000 Mathematics Subject Classification:** 05C75, 05C99.

## 1. Introduction

In this paper, we consider only finite, simple, undirected graphs with no loops and no multiple edges. For a graph  $G$ , we denote by  $V(G)$ ,  $E(G)$  and  $\Delta(G)$  the vertex set, the edge set and the maximum degree of  $G$ , respectively. For a vertex  $x$  of a graph  $G$ , the neighborhood of  $x$  in  $G$  is denoted by  $N_G(x)$ , and  $d_G(x) := |N_G(x)|$ .

For a subset  $S$  of  $V(G)$ , the subgraph in  $G$  induced by  $S$  is denoted by  $\langle S \rangle_G$ . For a subgraph  $H$  of  $G$ ,  $G - H = \langle V(G) - V(H) \rangle_G$ . For disjoint subsets  $S$  and  $T$  of  $V(G)$ , we let  $E_G(S, T)$  denote the set of edges of  $G$  joining a vertex in  $S$  and a vertex in  $T$ . When  $S$  or  $T$  consists of a single vertex, say  $S = \{x\}$  or  $T = \{y\}$ , we write  $E_G(x, T)$  or  $E_G(S, y)$  for  $E_G(S, T)$ . Let  $\overline{G}$  stand for the complement of  $G$ . For positive integers  $s, t$ , let  $r(s, t)$

be the Ramsey number, i.e., the smallest value of  $n$  for which every red-blue coloring of  $K_n$  yields a red  $K_s$  or a blue  $K_t$ . A graph  $G$  is said to be  $K_{1,t}$ -free if  $G$  contains no  $K_{1,t}$  as an induced subgraph. In particular, a graph  $G$  is said to be *claw-free* if  $G$  contains no  $K_{1,3}$  as an induced subgraph.

In this paper, we are concerned with a structure of graphs  $G$  for which both  $G$  and  $\overline{G}$  are  $K_{1,t}$ -free where  $t \geq 3$ . Our results are following.

**Theorem A.** *Let  $G$  be a graph with  $|V(G)| \geq 10$ . If both  $G$  and  $\overline{G}$  are claw-free, then  $\min\{\Delta(G), \Delta(\overline{G})\} \leq 2$ .*

**Theorem B.** *Let  $t$  be an integer with  $t \geq 4$ , and let  $G$  be a graph with  $|V(G)| \geq r(t^2 - t + 2, t^2 - t + 2)$ . If both  $G$  and  $\overline{G}$  are  $K_{1,t}$ -free, then  $\min\{\Delta(G), \Delta(\overline{G})\} \leq r(t - 1, t) - 1$ .*

In Theorem A, the bound on  $|V(G)|$  is best possible. To see this, we construct a graph  $G$  of order 9 such that  $\min\{\Delta(G), \Delta(\overline{G})\} > 2$ . Let  $C = v_1v_2 \dots v_8v_1$  be a cycle of length 8. Let  $v$  be a new vertex. Consider the graph  $G = (V(G), E(G))$  such that  $V(G) = V(C) \cup \{v\}$  and  $E(G) = E(C) \cup \{v_iv_j \mid 1 \leq i < j \leq 8, i+j \equiv 0 \pmod{4}\} \cup \{vv_{2l} \mid 1 \leq l \leq 4\}$ . Then  $|V(G)| = 9$ ,  $\min\{\Delta(G), \Delta(\overline{G})\} > 2$ , and both  $G$  and  $\overline{G}$  are claw-free (and isomorphic). Note that the converse of Theorem A is not true. To see this, consider  $G = K_3 \cup \overline{K_m}$  where  $m$  is a large positive integer. Then  $\min\{\Delta(G), \Delta(\overline{G})\} = \Delta(G) = 2$ . However, it is obvious that  $\overline{G}$  contains  $K_{1,3}$  as an induced subgraph. Avoiding this particular case, we obtain the following corollary.

**Corollary of Theorem A.** *Let  $G$  be a graph with  $|V(G)| \geq 10$ . Then the following statements are equivalent:*

- (i) *both  $G$  and  $\overline{G}$  are claw-free,*
- (ii) *either  $G$  or  $\overline{G}$  is a triangle-free graph of maximum degree at most 2.*

Alternatively, the statement (ii) can be formulated as follows.

- (ii) *either  $G$  or  $\overline{G}$  is a disjoint union of cycles of length  $l \geq 4$ , paths and isolated vertices.*

**Sketch of proof.**

(i) $\Rightarrow$ (ii). Theorem A implies that either  $G$  or  $\overline{G}$  (say,  $G$ ) has maximum degree at most 2. Then it is easy to see that  $G$  is also triangle-free: if  $\{x, y, z\} \subset V(G)$  induces a triangle in  $G$ , then this triangle is a component

of  $G$  since  $\Delta(G) \leq 2$ . Then for any vertex  $u \in V(G) \setminus \{x, y, z\}$  (which exists since  $|V(G)| \geq 10$ ), the set  $\{u, x, y, z\}$  induces a claw in  $\overline{G}$ , centered at  $u$ .

(ii) $\Leftrightarrow$ (i). Suppose that e.g.  $G$  is triangle-free with  $\Delta(G) \leq 2$ . Then  $G$  is claw-free since  $\Delta(G) \leq 2$  and  $\overline{G}$  is claw-free since  $G$  is triangle-free. For  $\overline{G}$  the proof is similar. ■

Theorem B is a similar result concerning graphs  $G$  for which both  $G$  and  $\overline{G}$  are  $K_{1,t}$ -free where  $t \geq 4$ . Now we show that there exists a graph  $G$  such that both  $G$  and  $\overline{G}$  are  $K_{1,t}$ -free and  $\min\{\Delta(G), \Delta(\overline{G})\} = r(t-1, t) - 1$ . Let  $R$  be a graph with  $|V(R)| = r(t-1, t) - 1$  such that  $R$  does not contain  $K_{t-1}$  or  $\overline{K}_t$  as an induced subgraph. Let  $v$  be a new vertex. Consider  $G = (R + v) \cup \overline{K_{|V(G)|-r(t-1,t)}}$  where  $|V(G)|$  is sufficiently large. Then  $G$  is a graph such that both  $G$  and  $\overline{G}$  are  $K_{1,t}$ -free and  $\min\{\Delta(G), \Delta(\overline{G})\} = r(t-1, t) - 1$ .

## 2. Proof of Theorem A

By contradiction, suppose that  $\Delta(G) \geq 3$  and  $\Delta(\overline{G}) \geq 3$ . Then by the assumption that both  $G$  and  $\overline{G}$  are claw-free,  $G$  contains a subgraph  $A$  such that  $A \cong K_3$  in  $G$ , and  $\overline{G}$  contains a subgraph  $B$  such that  $B \cong K_3$  in  $\overline{G}$ .

**Claim.** *Both  $G$  and  $\overline{G}$  do not contain a subgraph which is isomorphic to  $K_4$ .*

**Proof.** Suppose not. Then by symmetry, we may assume  $\overline{G} \supset K_4$ . Then  $G$  contains a subgraph  $S$  such that  $\langle V(S) \rangle_G \cong \overline{K}_4$ . Let  $V(S) = \{a, b, c, d\}$ . First suppose that  $G - S$  contains a subgraph  $T$  which is isomorphic to  $K_3$ . Let  $V(T) = \{e, f, g\}$ . Since  $\overline{G}$  is claw-free,  $E_G(x, V(T)) \neq \emptyset$  for every  $x \in V(S)$ . From  $|V(S)| = 4$ , there exists  $y \in V(T)$  such that  $|E_G(y, V(S))| \geq 2$ . By symmetry, we may assume  $ae, be \in E(G)$ . Since  $G$  is claw-free and  $\langle V(S) \rangle_G \cong \overline{K}_4$ , it follows that  $E_G(e, \{c, d\}) = \emptyset$ . By the assumption that  $G$  is claw-free,  $\langle \{e, a, b, f\} \rangle_G$  is not isomorphic to  $K_{1,3}$ . Hence by symmetry of the roles of  $a$  and  $b$ , we may assume  $af \in E(G)$ . Note that  $\langle \{a, e, f\} \rangle_G \cong K_3$ . Then by the assumption that  $\overline{G}$  is claw-free,  $E_G(c, \{a, e, f\}) \neq \emptyset$  and  $E_G(d, \{a, e, f\}) \neq \emptyset$ . This forces  $cf, df \in E(G)$ . Then  $\langle \{f, a, c, d\} \rangle_G \cong K_{1,3}$ . This is a contradiction. Hence it follows that  $G - S$  does not contain a triangle. Since  $G$  contains a triangle, we may assume that there exist  $u, v \in V(G - S)$  such that  $\langle \{a, u, v\} \rangle_G \cong K_3$ . Since  $\overline{G}$  is claw-free,  $|E_G(\{b, c, d\}, \{u, v\})| \geq 3$ . Then by the symmetry of the roles

of  $u$  and  $v$ , we may assume  $bu, cu \in E(G)$ . Then  $\langle \{u, a, b, c\} \rangle_G \cong K_{1,3}$ . This is a contradiction. ■

*Case 1.*  $V(A) \cap V(B) = \emptyset$ .

First suppose that there exist  $x, y \in V(B)$  such that  $N_G(x) \cap N_G(y) \cap V(A) \neq \emptyset$ . We may assume that  $V(A) = \{a, b, c\}$ ,  $V(B) = \{x, y, z\}$ , and  $ax, ay \in E(G)$ . Since  $\langle \{a, b, x, y\} \rangle_G \not\cong K_{1,3}$ , by the symmetry of the roles of  $x$  and  $y$ , we may assume  $bx \in E(G)$ . Since  $\langle \{a, b, x\} \rangle_G \cong K_3$ ,  $\langle \{a, x, y, z\} \rangle_G \not\cong K_{1,3}$  and  $\overline{G}$  is claw-free, it follows that  $bz \in E(G)$ . By the claim,  $xc \notin E(G)$ . Hence it follows from  $\langle \{b, c, x, z\} \rangle_G \not\cong K_{1,3}$  that  $cz \in E(G)$ . Since  $\langle \{b, c, z\} \rangle_G \cong K_3$  and both  $G$  and  $\overline{G}$  are claw-free, this forces  $cy \in E(G)$ . Hence by the claim, we have  $E_G(A, B) = \{ax, ay, bx, bz, cy, cz\}$ . Let  $v \in V(G - A - B)$ . Since  $\overline{G}$  is claw-free, it follows that  $E_G(v, V(A)) \neq \emptyset$ . By symmetry, we may assume  $va \in E(G)$ . Suppose that  $vc \in E(G)$ . Then it follows from the claim that  $E_G(v, \{b, y\}) = \emptyset$ . Then  $\langle a, b, y, v \rangle_G \cong K_{1,3}$ , a contradiction. Thus  $vc \notin E(G)$ . We can similarly obtain  $vb \notin E(G)$ . Since  $\langle \{a, v, x, c\} \rangle_G \not\cong K_{1,3}$  and  $\langle \{a, v, y, b\} \rangle_G \not\cong K_{1,3}$ , it follows that  $vx, vy \in E(G)$ . Since  $\overline{G}$  is claw-free,  $E_G(z, \{v, a, x\}) \neq \emptyset$ . This forces  $vz \in E(G)$ . Then  $\langle \{v, x, y, z\} \rangle_G \cong K_{1,3}$ , a contradiction. Thus we may assume that there exist no two vertices  $x, y \in V(B)$  such that  $N_G(x) \cap N_G(y) \cap V(A) \neq \emptyset$ . Since  $\overline{G}$  is claw-free, we may assume that  $V(A) = \{a, b, c\}$ ,  $V(B) = \{x, y, z\}$ , and  $E_G(V(A), V(B)) = \{ax, by, cz\}$ . Take  $v \in V(G - A - B)$ . Since  $\overline{G}$  is claw-free,  $E_G(v, V(A)) \neq \emptyset$ . By symmetry, we may assume  $av \in E(G)$ . By the claim,  $N_G(v) \not\subseteq V(A)$ . By symmetry, we may assume  $bv \notin E(G)$ . Since  $\langle \{a, v, x, b\} \rangle_G \not\cong K_{1,3}$ ,  $vx \in E(G)$ . Then  $\langle \{x, v, a\} \rangle_G \cong K_3$ . This forces  $vy, vz \in E(G)$  because  $\overline{G}$  is claw-free. Then  $\langle \{v, x, y, z\} \rangle_G \cong K_{1,3}$ . This is a contradiction.

*Case 2.*  $V(A) \cap V(B) \neq \emptyset$ .

We may assume that  $V(A) = \{a, b, c\}$  and  $V(B) = \{a, x, y\}$ . Since both  $G$  and  $\overline{G}$  are claw-free, it follows that either  $E_G(\{b, c\}, \{x, y\}) = \{bx, cy\}$  or  $E_G(\{b, c\}, \{x, y\}) = \{cx, by\}$ . By symmetry, we may assume  $E_G(\{b, c\}, \{x, y\}) = \{bx, cy\}$ . First suppose that there exist two vertices  $u, v \in V(G - A - B)$  such that either  $|E_G(b, \{u, v\})| = 2$  or  $|E_G(c, \{u, v\})| = 2$ . By symmetry, we may assume  $|E_G(c, \{u, v\})| = 2$ . Suppose that  $av \in E(G)$ . Then  $xv \in E(G)$  because  $\langle \{a, c, v\} \rangle_G \cong K_3$ . Also by the claim,  $vb \notin E(G)$ . Since  $\langle \{v\} \cup V(B) \rangle_G \not\cong K_{1,3}$ ,  $vy \notin E(G)$ . Then  $\langle \{c, v, y, b\} \rangle_G \cong K_{1,3}$ . This is a contradiction. Thus we have  $va \notin E(G)$ . By the symmetry of the roles of  $u$  and

$v$ , we can similarly have  $ua \notin E(G)$ . Then since  $\langle \{c, a, u, v\} \rangle_G \not\cong K_{1,3}$ , it follows that  $uv \in E(G)$ . Since  $\langle \{c, a, v, y\} \rangle_G \not\cong K_{1,3}$  and  $\langle \{c, a, u, y\} \rangle_G \not\cong K_{1,3}$ , it follows that  $uy, vy \in E(G)$ . Then  $\langle \{c, u, v, y\} \rangle_G \cong K_4$ , which contradicts the claim. Thus we may assume that there exist no two vertices  $u, v \in V(G - A - B)$  such that  $|E_G(b, \{u, v\})| = 2$  or  $|E_G(c, \{u, v\})| = 2$ . Then  $|E_G(a, V(G - A - B))| \geq 3$  because  $|V(G)| \geq 10$  (note that  $E_G(z, V(A)) \neq \emptyset$  for every  $z \in V(G - A - B)$  since  $\overline{G}$  is claw-free). Let  $u, v, w \in N_G(a) \cap V(G - A - B)$ . Since  $\langle \{a, u, v, w\} \rangle_G \not\cong K_{1,3}$ , we may assume  $uv \in E(G)$ . Since  $\langle \{a, u, v\} \rangle_G \cong K_3$  and  $\langle \{a, x, y\} \rangle_G \cong \overline{K_3}$ , either  $E_G(\{x, y\}, \{u, v\}) = \{vy, ux\}$  or  $E_G(\{x, y\}, \{u, v\}) = \{vx, uy\}$ . By symmetry, we may assume  $E_G(\{x, y\}, \{u, v\}) = \{vy, ux\}$ . If  $\langle \{c, a, v\} \rangle_G \cong K_3$ , then  $xv \in E(G)$ , which implies  $\langle \{v, a, x, y\} \rangle_G \cong K_{1,3}$ , a contradiction. Hence  $vc \notin E(G)$ . We can similarly have  $ub \notin E(G)$ . Suppose that  $E_G(w, \{u, v\}) = \emptyset$ . Then since  $\langle \{a, b, u, w\} \rangle_G \not\cong K_{1,3}$  and  $\langle \{a, c, v, w\} \rangle_G \not\cong K_{1,3}$ , this forces  $wb, wc \in E(G)$ , which contradicts the claim. Thus we may assume  $E_G(w, \{u, v\}) \neq \emptyset$ . By the claim, note that  $|E_G(w, \{u, v\})| \leq 1$ . By symmetry, we may assume  $E_G(w, \{u, v\}) = \{wv\}$ . Then  $xw \in E(G)$  because  $\langle \{a, w, v\} \rangle_G \cong K_3$ . Since  $\langle \{a, b, u, w\} \rangle_G \not\cong K_{1,3}$ , this forces  $bw \in E(G)$ . Then we have  $yw \in E(G)$  because  $\langle \{a, b, w\} \rangle_G \cong K_3$ . Then  $\langle \{w, a, x, y\} \rangle_G \cong K_{1,3}$ . This is a contradiction. This completes the proof of Theorem A. ■

### 3. Proof of Theorem B

By contradiction, suppose that  $\Delta(G) \geq r(t-1, t)$  and  $\Delta(\overline{G}) \geq r(t-1, t)$ . Then both  $G$  and  $\overline{G}$  contain  $K_t$  because both  $G$  and  $\overline{G}$  are  $K_{1,t}$ -free. Let  $A$  be a subgraph of  $G$  such that  $A \cong K_t$ . Since  $|V(G)| \geq r(t^2 - t + 2, t^2 - t + 2)$ , by symmetry, we may assume that  $G$  contains  $\overline{K_{t^2-t+2}}$  as an induced subgraph. Hence there exists a subgraph  $H$  of  $G$  such that  $\langle V(H) \rangle_G \cong \overline{K_{t^2-t+1}}$  and  $V(A) \cap V(H) = \emptyset$ . Since  $\overline{G}$  is  $K_{1,t}$ -free,  $E_G(x, V(A)) \neq \emptyset$  for every  $x \in V(H)$ . Since  $|V(H)| = t(t-1) + 1$ , there exists a  $v \in V(A)$  such that  $|E_G(v, V(H))| \geq t$ . This implies that  $\langle \{v\} \cup V(H) \rangle_G$  contains  $K_{1,t}$  as an induced subgraph. This is a contradiction. This completes the proof of Theorem B. ■

**Remark.** Theorem A implies as an immediate corollary that the graph property “both  $G$  and  $\overline{G}$  are claw-free” is stable under the closure for claw-free graphs, i.e., if  $G$  has the property, then  $cl(G)$  has the property as well (see e.g. the survey paper [1]).

## References

- [1] H.J. Broersma, Z. Ryjáček and I. Schiermeyer, *Closure concepts — a survey*, Graphs and Combin. **16** (2000) 17–48.

Received 23 November 2003

Revised 6 November 2004