

## ON RAMSEY $(K_{1,2}, C_4)$ -MINIMAL GRAPHS

TOMÁŠ VETRÍK

*School of Mathematical Sciences*  
*University of KwaZulu-Natal*  
*Durban, South Africa*

**e-mail:** tomas.vetrik@gmail.com

LYRA YULIANTI AND EDY TRI BASKORO

*Combinatorial Mathematics Research Division*  
*Faculty of Mathematics and Natural Sciences*  
*Institut Teknologi Bandung*  
*Bandung, Indonesia*

**e-mail:** lyra@students.itb.ac.id

**e-mail:** ebaskoro@math.itb.ac.id

### Abstract

For graphs  $F$ ,  $G$  and  $H$ , we write  $F \rightarrow (G, H)$  to mean that any red-blue coloring of the edges of  $F$  contains a red copy of  $G$  or a blue copy of  $H$ . The graph  $F$  is Ramsey  $(G, H)$ -minimal if  $F \rightarrow (G, H)$  but  $F^* \not\rightarrow (G, H)$  for any proper subgraph  $F^* \subset F$ . We present an infinite family of Ramsey  $(K_{1,2}, C_4)$ -minimal graphs of any diameter  $\geq 4$ .

**Keywords:** Ramsey-minimal graph, edge coloring, diameter of a graph.

**2010 Mathematics Subject Classification:** 05C55, 05D10.

### 1. INTRODUCTION

All graphs considered in this paper are finite, undirected, without loops and multiple edges. Let  $G$  be a graph with the vertex set  $V(G)$  and the edge set  $E(G)$ . The *distance*  $d_G(u, v)$  between two vertices  $u$  and  $v$  in a graph  $G$  is the length of the shortest path connecting them. The *eccentricity* of a vertex  $u$  is the greatest distance between  $u$  and any other vertex in  $G$ . The *diameter*

of a connected graph  $G$  is the maximum distance between two vertices in  $G$ . If  $G$  contains vertices  $v_1, v_2, v_3, v_4$  and edges  $v_1v_2, v_2v_3, v_3v_4, v_4v_1, v_1v_3$ , we say that the edge  $v_1v_3$  lies inside 4-cycle  $v_1v_2v_3v_4$ .

Let  $F, G$  and  $H$  be graphs. We say that  $F$  contains  $G$  if  $F$  contains a subgraph isomorphic to  $G$ . We write  $F \rightarrow (G, H)$  if whenever each edge of  $F$  is colored either red or blue, then  $F$  contains a red copy of  $G$  or a blue copy of  $H$ . A graph  $F$  is Ramsey  $(G, H)$ -minimal if  $F \rightarrow (G, H)$  but  $F^* \not\rightarrow (G, H)$  for any proper subgraph  $F^* \subset F$ . The class of all Ramsey  $(G, H)$ -minimal graphs is denoted by  $\mathcal{R}(G, H)$ .

Numerous papers study the problem of determining the set  $\mathcal{R}(G, H)$ . Burr, Erdős and Lovász [5] showed that  $\mathcal{R}(K_{1,2}, K_{1,2}) = \{K_{1,3}, C_{2n+1}\}$  where  $n \geq 1$ . Later, Burr *et al.* [4] proved that if  $m, n$  are odd, then  $\mathcal{R}(K_{1,m}, K_{1,n}) = \{K_{1,m+n-1}\}$ . All graphs belonging to  $\mathcal{R}(2K_2, K_{1,n})$  for  $n \geq 3$  were presented by Mengersen and Oeckermann [7]. Borowiecki, Hałuszczak and Sidorowicz [2] determined the class  $\mathcal{R}(K_{1,2}, K_{1,n})$  for  $n \geq 3$ .

Luczak [6] proved that if  $G$  is a forest other than a matching and  $H$  is a graph containing at least one cycle, then  $\mathcal{R}(G, H)$  is infinite. It follows that the set  $\mathcal{R}(K_{1,2}, C_n)$  is infinite for any  $n \geq 3$ . Borowiecki, Schiermeyer and Sidorowicz [3] found all graphs in  $\mathcal{R}(K_{1,2}, C_3)$ . Recently, Baskoro, Yulianti and Assiyatun [1] gave a family of graphs belonging to  $\mathcal{R}(K_{1,2}, C_4)$ , where an infinite family of Ramsey  $(K_{1,2}, C_4)$ -minimal graphs was stated only for diameter 2. We present an infinite class of Ramsey  $(K_{1,2}, C_4)$ -minimal graphs for any diameter  $\geq 4$ .

## 2. GRAPHS OF DIAMETER 4

We define some classes of graphs. Let  $t \geq 6$  be an even integer. Let  $G(t)$  be a graph with the vertex set  $V(G(t)) = \{v, v_1, v_2, \dots, v_t = v_0\}$  and with the edge set  $E(G(t)) = \{vv_{2i} : i = 1, 2, \dots, \frac{t}{2}\} \cup \{v_jv_{j+1} : j = 0, 1, \dots, t-1\}$ .

Let  $A_1(t)$  be a graph with  $V(A_1(t)) = V(G(t)) \cup \{v', v'_0\}$  and  $E(A_1(t)) = E(G(t)) \cup \{vv_1, vv'_0, v_0v'_0, v'v_0, v'v_1, v_2v_4\}$ .

Let  $A_2(t)$  be a graph with  $V(A_2(t)) = V(G(t)) \cup \{v'_1, v'_p\}$  for odd  $p \in \{3, 5, \dots, t-1\}$  and  $E(A_2(t)) = E(G(t)) \cup \{v_0v'_1, v'_1v_2, v_{p-1}v'_p, v'_pv_{p+1}\}$ .

Let  $A_3(t)$  be a graph with  $V(A_3(t)) = V(G(t)) \cup \{v'_1, v'_2\}$  and  $E(A_3(t)) = E(G(t)) \cup \{vv_3, vv'_2, v_2v'_2, v_0v'_1, v'_1v_2\}$ .

We show that  $A_1(t), A_2(t)$  and  $A_3(t)$  are Ramsey  $(K_{1,2}, C_4)$ -minimal graphs.

**Assertion 1.**  $A_1(t) \in \mathcal{R}(K_{1,2}, C_4)$ .

**Proof.** First we prove that  $A_1(t) \rightarrow (K_{1,2}, C_4)$ . Consider any red-blue coloring of the edges of  $A_1(t)$ . Suppose that there is no red copy of  $K_{1,2}$  in the coloring. Since the edges  $vv_0, vv_1, vv_2, v_0v_1$  and  $v_2v_4$  lie inside 4-cycles, we can not color them by red, because otherwise, we would have blue copies of  $C_4$  in our coloring. We must color by red the edge  $v_1v_2$  to avoid blue 4-cycle  $vv_0v_1v_2$ . Next, we must color by red the edge  $v'v_0$  to avoid blue 4-cycle  $vv_0v'v_1$  and the edge  $vv'_0$  to avoid blue 4-cycle  $vv'_0v_0v_1$ . Then all the edges  $vv_i, i = 0, 2, \dots, t - 2$  must be blue. It follows that to avoid blue 4-cycles  $vv_jv_{j+1}v_{j+2}, j = 2, 4, \dots, t - 4$ , the edges  $v_{j+1}v_{j+2}$  must be red and  $v_jv_{j+1}$  are blue. But since  $v_{t-3}v_{t-2}$  and  $v'v_0$  are red, we are not able to avoid blue 4-cycle  $vv_{t-2}v_{t-1}v_0$  which means that  $A_1(t) \rightarrow (K_{1,2}, C_4)$ .

Now let us show that  $A_1^*(t) \not\rightarrow (K_{1,2}, C_4)$  for the graph  $A_1^*(t) \simeq A_1(t) \setminus \{e\}$ , where  $e$  is any fixed edge of  $A_1(t)$ . Let  $e = v_lv_{l+1}, l = 2, 3, \dots, t - 1$ . We can color by red the edges  $vv'_0, v'v_0$  and  $v_iv_{i+1}$ , where  $i = 1, 3, \dots, l - 1; l + 2, l + 4, \dots, t - 2$  if  $l$  is even, and  $i = 1, 3, \dots, l - 2; l + 1, l + 3, \dots, t - 2$  if  $l$  is odd. We color by blue all the edges of  $A_1^*(t)$  that are not colored by red.

If  $e = vv_l, l = 2, 4, \dots, t$ , color by red the edges  $vv'_0, v'v_0, v_1v_2$  and  $v_iv_{i+1}, i = 3, 5, \dots, l - 3; l + 2, l + 4, \dots, t - 2$ . If  $e = vv'_0, v_0v'_0$  or  $v_0v_1$ , the edges colored by red are  $vv_0$  and  $v_iv_{i+1}$ , where  $i = 1, 3, \dots, t - 3$ . If  $e = vv_1, v_1v_2$  or  $v_2v_4$ , we can color by red  $v'v_1, v_0v'_0, vv_2$  and  $v_iv_{i+1}, i = 4, 6, \dots, t - 2$ . Finally, if  $e = v'v_0$  or  $v'v_1$ , we color by red  $vv_4, v_0v_1$  and  $v_iv_{i+1}, i = 6, 8, \dots, t - 2$ . The other edges will be colored by blue. These colorings of  $A_1^*(t)$  contain neither a red copy of  $K_{1,2}$  nor a blue copy if  $C_4$ . The proof is complete. ■

**Assertion 2.**  $A_2(t) \in \mathcal{R}(K_{1,2}, C_4)$ .

**Proof.** Let us show that  $A_2(t) \rightarrow (K_{1,2}, C_4)$ . We consider any red-blue coloring of the edges of  $A_2(t)$  such that there is no red copy of  $K_{1,2}$  in the coloring. In order to avoid blue 4-cycles containing at least one of the vertices  $v_1, v'_1, v_p$  or  $v'_p$ , we must color by red one of the edges  $v_iv, v_iv_1, v_iv'_1$  for  $i = 0, 2$  and one of the edges  $v_jv, v_jv_p, v_jv'_p$  for  $j = p - 1, p + 1$ . Note that if  $p = 3$  or  $p = t - 1$ , we must color by red the edge  $vv_2$  or  $vv_0$ . There can be at most one red edge  $vv_i, i \in \{2, 4, \dots, t\}$  in our coloring. It can be seen that if all the edges  $vv_i, i = 2, 4, \dots, p - 1$  are blue, we can not avoid blue 4-cycle  $vv_jv_{j+1}v_{j+2}$  for some  $j \in \{2, 4, \dots, p - 3\}$ , and if all the edges

$vv_i, i = p+1, p+3, \dots, t$  are blue, it is not possible to avoid blue 4-cycle  $vv_jv_{j+1}v_{j+2}$  for a  $j \in \{p+1, p+3, \dots, t-2\}$ . Therefore,  $A_2(t) \rightarrow (K_{1,2}, C_4)$ .

To prove the minimality of  $A_2(t)$ , consider the graph  $A_2^*(t) \simeq A_2(t) \setminus \{e\}$  for any fixed edge  $e \in E(A_2(t))$ . Let  $e = v_l v_{l+1}, l = 0, 1, \dots, p$ . We can color by red the edges  $vv_0, v'_p v_{p+1}, v_i v_{i+1}, i = p+2, p+4, \dots, t-3$  and  $v_j v_{j+1}$ , where  $j = 1, 3, \dots, l-1; l+2, l+4, \dots, p-1$  if  $l$  is even, and  $j = 1, 3, \dots, l-2; l+1, l+3, \dots, p-1$  if  $l$  is odd. If  $e = vv_l, l = 2, 4, \dots, p+1$ , the edges colored by red are  $vv_0, v'_1 v_2, v'_p v_{p+1}$  and  $v_i v_{i+1}, i = 3, 5, \dots, l-3; l+2, l+4, \dots, p-1; p+2, p+4, \dots, t-3$ . The rest of the edges of  $A_2^*(t)$  will be colored by blue. There is no red copy of  $K_{1,2}$  and no blue copy of  $C_4$  in these colorings. The cases  $e = v_0 v'_1, v'_1 v_2, v_{p-1} v'_p, v'_p v_{p+1}, v_j v_{j+1}, j = p+1, p+2, \dots, t-1$  or  $e = vv_i, i = p+3, p+5, \dots, t$  are similar. ■

**Assertion 3.**  $A_3(t) \in \mathcal{R}(K_{1,2}, C_4)$ .

**Proof.** We show that  $A_3(t) \rightarrow (K_{1,2}, C_4)$ . Let us consider any red-blue coloring of  $A_3(t)$ . Assume there is no red  $K_{1,2}$  in the coloring. We can not color by red the edges  $vv_2$  and  $vv_3$ , because they lie inside 4-cycles  $vv'_2 v_2 v_3$  and  $vv_2 v_3 v_4$ . We also can not color by red the edges  $v_2 v'_2$  and  $v_2 v_3$ , because then, we would not be able to avoid blue 4-cycle  $vv_0 v_1 v_2, vv_0 v'_1 v_2$  or  $v_0 v_1 v_2 v'_1$ . It follows that to avoid blue 4-cycle  $vv'_2 v_2 v_3$ , we must color by red the edge  $vv'_2$ . Then the edges  $vv_i, i = 2, 4, \dots, t$  must be blue. Consequently, if we want to avoid blue cycles  $vv_0 v_1 v_2$  and  $vv_0 v'_1 v_2$ , we must color by red either the edges  $v_0 v_1, v'_1 v_2$  or the edges  $v_0 v'_1, v_1 v_2$ . The edges  $v_j v_{j+1}, j = 2, 3, \dots, t-3$  must be colored alternately by blue and red. It follows that we can not avoid blue 4-cycle  $vv_{t-2} v_{t-1} v_t$ . Hence,  $A_3(t) \rightarrow (K_{1,2}, C_4)$ .

In order to prove the minimality of  $A_3(t)$  we consider  $A_3^*(t) \simeq A_3(t) \setminus \{e\}$ , where  $e$  is any fixed edge of  $A_3(t)$ . Let  $e = v_l v_{l+1}, l = 0, 1, \dots, t-1$ . We can color by red the edges  $vv'_2, v_0 v'_1$  and  $v_i v_{i+1}$ , where  $i = 1, 3, \dots, l-1; l+2, l+4, \dots, t-2$  if  $l$  is even (where  $i = 1, 3, \dots, l-2; l+1, l+3, \dots, t-2$  if  $l$  is odd). If  $e = vv_l, l = 2, 4, \dots, t$ , the edges colored by red are  $vv'_2, v_0 v'_1, v_1 v_2$  and  $v_i v_{i+1}, i = 1, 3, \dots, l-3; l+2, l+4, \dots, t-2$ . If  $e = vv_3, vv'_2$  or  $v_2 v'_2$ , color by red  $vv_2, v_0 v'_1$  and  $v_i v_{i+1}$ , where  $i = 4, 6, \dots, t-2$ , and if  $e = v_0 v'_1$  or  $v'_1 v_2$ , color by red  $vv_0, v_2 v'_2$  and  $v_i v_{i+1}, i = 3, 5, \dots, t-3$ . The other edges will be colored by blue. The colorings of  $A_3^*(t)$  contain neither a red  $K_{1,2}$  nor a blue  $C_4$ . This finishes the proof. ■

It is easy to verify that the graphs  $A_i(t), i = 1, 2, 3$  have diameter 4 for  $t \geq 8$ , and 3 if  $t = 6$ .

3. AUXILIARY RESULTS

Let us introduce Definitions 1 and 2.

**Definition 1.** Let  $F$  be a graph with  $U \subset V(F)$ . For any given graphs  $G$  and  $H$ , provided that the vertices in  $U$  are not incident to red edges, we write  $F \rightarrow (G(U), H)$  to mean that any red-blue coloring of the edges of  $F$  contains a red copy of  $G$  or a blue copy of  $H$ .

**Definition 2.** Let  $U_0 \subset V(F)$  where  $|U_0| = p$ . For  $i \in \{0, 1, \dots, p - 1\}$  a graph  $F$  is Ramsey  $(G(U_0)_i, H)$ -minimal if

- (i)  $F \rightarrow (G(U_i), H)$ , where  $U_i$  is any subset of  $U_0$  such that  $|U_i| = p - i$ ,
- (ii)  $F^* \not\rightarrow (G(U_i), H)$  for any proper subgraph  $F^* \subset F$ ,
- (iii)  $F \not\rightarrow (G(U_{i+1}), H)$ , where  $U_{i+1}$  is any subset of  $U_i$  such that  $|U_{i+1}| = p - i - 1$ .

Vertices in  $U_0$  will be called *roots* of  $F$  and the class of all Ramsey  $(G(U_0)_i, H)$ -minimal graphs will be denoted by  $\mathcal{R}(G(U_0)_i, H)$ .

If  $F$  is Ramsey  $(G(U_0)_0, H)$ -minimal, we write  $F \in \mathcal{R}(G(U_0), H)$ . Particularly, for  $U_0 = \emptyset$ ,  $F$  is a Ramsey  $(G, H)$ -minimal graph.

We need to define the following families of graphs:

$L_1(t)$  is a graph with  $V(L_1(t)) = V(G(t)) \cup \{v'\}$  and  $E(L_1(t)) = E(G(t)) \cup \{vv_1, v'v_0, v'v_1, v_2v_4\}$ . Let us remind that  $G(t)$  is defined for an even integer  $t \geq 6$ .

$L_2(t)$  is a graph with  $V(L_2(t)) = V(G(t)) \cup \{v'_1\}$  and  $E(L_2(t)) = E(G(t)) \cup \{v_0v'_1, v'_1v_2\}$ .

$L_3(t)$  is a graph with  $V(L_3(t)) = V(G(t)) \cup \{v'_0\}$  and  $E(L_3(t)) = E(G(t)) \cup \{vv_1, vv'_0, v_0v'_0\}$ .

$M_2(t) = G(t)$  and  $M_3(t)$  is a graph with  $V(M_3(t)) = V(G(t))$  and  $E(M_3(t)) = E(G(t)) \setminus \{v_0v_1, v_1v_2\} \cup \{vv_1, v_1v_4, vv_5\}$ .

Let  $s \geq 5$  be odd.  $M_1(s)$  is a graph with the vertex set  $V(M_1(s)) = \{v, v_1, v_2, \dots, v_s = v_0\}$  and with the edge set  $E(M_1(s)) = \{vv_i, i = 1, 2, \dots, s\} \cup \{v_jv_{j+1}, j = 1, 2, \dots, s - 1\}$ .

We prove some lemmas characterizing the graphs defined above.

**Lemma 1.** (i) Let  $t \geq 8$  and  $p \in \{6, 8, \dots, t - 2\}$ . Then  $L_1(t) \in \mathcal{R}(K_{1,2}(v_p), C_4)$ .

(ii) Let  $t \geq 10$  and  $r, s \in \{6, 8, \dots, t - 2\}$ ,  $r \neq s$ . Then  $L_1(t) \in \mathcal{R}(K_{1,2}(v_r, v_s)_1, C_4)$ .

**Proof.** (i) First we show that  $L_1(t) \rightarrow (K_{1,2}(v_p), C_4)$  for even integers  $t, p$ , where  $t \geq 8$  and  $p \in \{6, 8, \dots, t-2\}$ . Provided that there are no red edges incident to the vertex  $v_p$ , let us consider any red-blue coloring of the edges of  $L_1(t)$  such that we have no red copy of  $K_{1,2}$  in the coloring. Since the edges  $vv_1, vv_2, v_0v_1$  and  $v_2v_4$  lie inside 4-cycles, we can not color them by red, because then, we would have blue copies of  $C_4$  in our coloring. We must color by red one of the edges  $vv_0, v_1v_2$  and one of the edges  $vv_4, v_1v_2$  to avoid blue 4-cycles  $vv_0v_1v_2$  and  $vv_1v_2v_4$ , which means that  $v_1v_2$  must be red in any case. Consequently, we color by red one of the edges  $vv_4, v_3v_4$  and one of the edges  $vv_0, v'v_0$  to avoid blue 4-cycles  $vv_2v_3v_4$  and  $vv_0v'v_1$ .

Since there can be at most one red edge  $vv_i, i \in \{4, 6, \dots, t\}, i \neq p$ , without lose of generality we can assume that all the edges  $vv_j, j = 4, 6, \dots, p-2$  are blue. In order to avoid blue 4-cycles  $vv_jv_{j+1}v_{j+2}, j = 2, 4, \dots, p-4$ , we must color the edges  $v_{j+1}v_{j+2}$  by red. Clearly, the edges  $v_jv_{j+1}$  are blue. Then, since  $v_{p-3}v_{p-2}$  is red and no red edge can be incident to  $v_p$ , we have blue 4-cycle  $vv_{p-2}v_{p-1}v_p$  in our coloring. Hence,  $L_1(t) \rightarrow (K_{1,2}(v_p), C_4)$ .

Now we prove that  $L_1^*(t) \rightarrow (K_{1,2}(v_p), C_4)$ , where  $L_1^*(t) \simeq L_1(t) \setminus \{e\}$  for any fixed edge  $e \in E(L_1(t))$ . Let  $e = v_lv_{l+1}, l = 2, 3, \dots, p-1$ . The edges colored by red are  $vv_0, v_iv_{i+1}, i = p+1, p+3, \dots, t-3$  and  $v_jv_{j+1}$ , where  $j = 1, 3, \dots, l-1; l+2, l+4, \dots, p-2$  if  $l$  is even, and  $j = 1, 3, \dots, l-2; l+1, l+3, \dots, p-2$  if  $l$  is odd.

If  $e = vv_l, l = 2, 4, \dots, p$ , we can color by red the edges  $vv_0, v_1v_2, v_iv_{i+1}, i = 3, 5, \dots, l-3; l+2, l+4, \dots, p-2; p+1, p+3, \dots, t-3$ . If  $e = vv_0, vv_1, v'v_0$  or  $v'v_1$ , color by red the edges  $vv_{p-2}$  and  $v_iv_{i+1}$ , where  $i = 1, 3, \dots, p-5; p+1, p+3, \dots, t-1$ . If  $e = v_0v_1$  or  $v_1v_2$ , the edges colored by red are  $vv_1$  and  $v_iv_{i+1}, i = 2, 4, \dots, p-2; p+1, p+3, \dots, t-1$ . If  $e = v_2v_4$ , we color by red  $v'v_1, vv_2$  and  $v_iv_{i+1}, i = 4, 6, \dots, p-2; p+1, p+3, \dots, t-1$ . The rest of the edges will be colored by blue. If  $e = v_lv_{l+1}, l = p, p+1, \dots, t-1$  or  $e = vv_k, k = p+2, p+4, \dots, t-2$ , we can analogously show that there exists a red-blue coloring of  $L_1^*(t)$  containing neither a red  $K_{1,2}$  nor a blue  $C_4$  such that there is no red edge incident to the vertex  $v_p$ .

Clearly,  $L_1(t) \rightarrow (K_{1,2}, C_4)$ , because  $L_1(t) \subset A_1(t)$ . Hence,  $L_1(t) \in \mathcal{R}(K_{1,2}(v_p), C_4)$ .

(ii) From the proof of part (i) we get  $L_1(t) \rightarrow (K_{1,2}(v_p), C_4)$  for  $p \in \{6, 8, \dots, t-2\}$ ,  $L_1^*(t) \rightarrow (K_{1,2}(v_p), C_4)$  for  $L_1^*(t) \simeq L_1(t) \setminus \{e\}$ , where  $e$  is any fixed edge of  $L_1(t)$ , and  $L_1(t) \rightarrow (K_{1,2}, C_4)$ . This shows that for  $t \geq 10$  one has  $L_1(t) \in \mathcal{R}(K_{1,2}(v_r, v_s)_1, C_4)$ , where  $r, s \in \{6, 8, \dots, t-2\}, r \neq s$ . The proof is complete. ■

- Lemma 2.** (i) Let  $t \geq 6$  and  $p \in \{4, 6, \dots, t-2\}$ . Then  $L_2(t) \in \mathcal{R}(K_{1,2}(v_p), C_4)$ .  
 (ii) Let  $t \geq 8$  and  $r, s \in \{4, 6, \dots, t-2\}$ ,  $r \neq s$ . Then  $L_2(t) \in \mathcal{R}(K_{1,2}(v_r, v_s)_1, C_4)$ .

**Proof.** (i) We prove that  $L_2(t) \rightarrow (K_{1,2}(v_p), C_4)$ . Consider any red-blue coloring of the edges of  $L_2(t)$  such that there is no red edge incident to the vertex  $v_p$ . Assume that we have no red  $K_{1,2}$  in the coloring. We must color by red one of the edges  $v_i v, v_i v_1, v_i v'_1$  for  $i = 0, 2$  to avoid blue 4-cycles containing at least one of the vertices  $v_1, v'_1$ . Note that there can be at most one red edge  $vv_i, i \in \{2, 4, \dots, t\}, i \neq p$  in our coloring. It is easy to show that if all the edges  $vv_i, i = 2, 4, \dots, p-2$  are blue, we are not able to avoid blue 4-cycle  $vv_j v_{j+1} v_{j+2}$  for some  $j \in \{2, 4, \dots, p-2\}$ , and if  $vv_i, i = p+2, p+4, \dots, t$  are blue, we can not avoid blue 4-cycle  $vv_j v_{j+1} v_{j+2}$  for a  $j \in \{p, p+2, \dots, t-2\}$ .  $L_2(t) \rightarrow (K_{1,2}(v_p), C_4)$ .

Consider  $L_2^*(t) \simeq L_2(t) \setminus \{e\}$  for any fixed edge  $e \in E(L_2(t))$ . We show that  $L_2^*(t) \not\rightarrow (K_{1,2}(v_p), C_4)$ . Let  $e = v_l v_{l+1}, l = 0, 1, \dots, p-1$ . We can color by red the edges  $vv_0, v_i v_{i+1}, i = p+1, p+3, \dots, t-3$  and the edges  $v_j v_{j+1}$ , where  $j = 1, 3, \dots, l-1; l+2, l+4, \dots, p-2$  if  $l$  is even, and  $j = 1, 3, \dots, l-2; l+1, l+3, \dots, p-2$  if  $l$  is odd. If  $e = vv_l, l = 2, 4, \dots, p$ , the edges colored by red are  $vv_0, v_1 v_2, v_i v_{i+1}, i = 3, 5, \dots, l-3; l+2, l+4, \dots, p-2; p+1, p+3, \dots, t-3$ . The other edges are colored by blue. The cases  $e = v_0 v'_1, v'_1 v_2, v_l v_{l+1}, l = p, p+1, \dots, t-1$  and  $e = vv_k, k = p+2, p+4, \dots, t$  are similar.

Finally, since  $L_2(t) \subset A_2(t)$ , it is evident that  $L_2(t) \not\rightarrow (K_{1,2}, C_4)$ .

- (ii) The proof follows from the previous part. ■

- Lemma 3.** (i) Let  $t \geq 6$  and  $p = 0$  or  $t-2$ . Then  $L_3(t) \in \mathcal{R}(K_{1,2}(v_p), C_4)$ .  
 (ii) Let  $t \geq 6$ . Then  $L_3(t) \in \mathcal{R}(K_{1,2}(v_0, v_{t-2})_1, C_4)$ .

The proof is analogous to the proofs of Lemma 1 and Lemma 2.

**Lemma 4.** Let  $s \geq 5$ . Then  $M_1(s) \in \mathcal{R}(K_{1,2}(v_1, v_s), C_4)$ .

**Proof.** Let us show that  $M_1(s) \rightarrow (K_{1,2}(v_1, v_s), C_4)$ . Provided that the vertices  $v_1, v_s$  are not incident to red edges, we consider any red-blue coloring of  $M_1(s)$  such that there is no red copy of  $K_{1,2}$  in the coloring. If we color by red some edge  $vv_i, i \in \{2, 3, \dots, s-1\}$ , we have blue 4-cycle  $vv_{i-1} v_i v_{i+1}$ . Therefore, all the edges  $vv_i, i = 1, 2, \dots, s$  must be blue. In order to avoid

blue 4-cycles  $vv_{j-1}v_jv_{j+1}$  and  $vv_jv_{j+1}v_{j+2}$ ,  $j = 2, 4, \dots, s - 3$ , the edges  $v_jv_{j+1}$  must be red. Then we are not able to avoid blue 4-cycle  $vv_{s-2}v_{s-1}v_s$ .

We prove that  $M_1^*(s) \not\rightarrow (K_{1,2}(v_1, v_s), C_4)$ , where  $M_1^*(s) \simeq M_1(s) \setminus \{e\}$  for any fixed edge  $e \in E(M_1(s))$ . Let  $e = v_lv_{l+1}$ ,  $l = 1, 2, \dots, s - 1$ . We can color by red the edges  $v_iv_{i+1}$ , where  $i = 2, 4, \dots, l - 2; l + 1, l + 3, \dots, s - 2$  if  $l$  is even, and  $i = 2, 4, \dots, l - 1; l + 2, l + 4, \dots, s - 2$  if  $l$  is odd. Let  $e = vv_l$ ,  $l = 3, 4, \dots, s$ . The edges colored by red are  $vv_{l-1}$  and  $v_iv_{i+1}$ , where  $i = 2, 4, \dots, l - 4, l + 1, l + 3, \dots, s - 2$  if  $l$  is even, and  $i = 2, 4, \dots, l - 3, l + 2, l + 4, \dots, s - 2$  if  $l$  is odd. We color by blue all the edges of  $M_1^*(s)$  that are not colored by red. The cases  $e = vv_1$  or  $vv_2$  can be handled similarly.

Finally,  $M_1(s) \not\rightarrow (K_{1,2}(v_p), C_4)$  for  $p = 1$  (for  $p = s$ ), since there exists a red-blue coloring of  $M_1(s)$  containing neither a red  $K_{1,2}$  nor a blue  $C_4$  such that there is no red edge incident to  $v_p$ . It is enough to color by red the edges  $v_iv_{i+1}$ , where  $i = 2, 4, \dots, s - 1$  (where  $i = 1, 3, \dots, s - 2$ ) and color by blue the rest of the edges. This finishes the proof. ■

**Lemma 5.** *Let  $t \geq 6$ . Then  $M_3(t) \in \mathcal{R}(K_{1,2}(v_0, v_2), C_4)$ .*

**Proof.** Let us consider any red-blue coloring of  $M_3(t)$  such that the vertices  $v_0, v_2$  are not incident to any red edges. We show that  $M_3(t) \rightarrow (K_{1,2}(v_0, v_2), C_4)$ . Suppose that we have no red  $K_{1,2}$  in the coloring. We can not color by red the edges  $vv_4$  and  $vv_5$ , because they lie inside 4-cycles  $vv_1v_4v_5$  and  $vv_4v_5v_6$ . It follows that we must color by red the edge  $v_3v_4$  to avoid blue cycle  $vv_2v_3v_4$ , and the edge  $vv_1$  to avoid blue cycle  $vv_1v_4v_5$ . But then, it is not possible to avoid blue 4-cycle  $vv_jv_{j+1}v_{j+2}$  for some  $j \in \{4, 6, \dots, t - 2\}$ , which shows that  $M_3(t) \rightarrow (K_{1,2}(v_0, v_2), C_4)$ .

Now consider the graph  $M_3^*(t) \simeq M_3(t) \setminus \{e\}$ , where  $e$  is any fixed edge of  $M_3(t)$ . Let us prove that  $M_3^*(t) \not\rightarrow (K_{1,2}(v_0, v_2), C_4)$ . Let  $e = v_lv_{l+1}$ ,  $l = 2, 3, \dots, t - 1$ . We can color by red the edges  $vv_1$  and  $v_iv_{i+1}$ , where  $i = 3, 5, \dots, l - 1; l + 2, l + 4, \dots, t - 2$  if  $l$  is even, and  $i = 3, 5, \dots, l - 2; l + 1, l + 3, \dots, t - 2$  if  $l$  is odd. If  $e = vv_l$ ,  $l = 2, 4, \dots, t$ , the edges colored by red are  $vv_1$  and  $v_iv_{i+1}$ , where  $i = 3, 5, \dots, l - 3; l + 2, l + 4, \dots, t - 2$ . If  $e = vv_1, vv_5$  or  $v_1v_4$ , we color by red the edges  $vv_4$  and  $v_iv_{i+1}$ ,  $i = 6, 8, \dots, t - 2$ . The rest of the edges will be colored by blue. The colorings of  $M_3^*(t)$  contain neither a red copy of  $K_{1,2}$  nor a blue copy of  $C_4$ .

In order to show that  $M_3(t) \not\rightarrow (K_{1,2}(v_p), C_4)$  for  $p = 0$  (for  $p = 2$ ) it suffices to color by red the edges  $v_iv_{i+1}$ ,  $i = 2, 4, \dots, t - 2$  (the edges  $vv_1$  and  $v_iv_{i+1}$ ,  $i = 3, 5, \dots, t - 1$ ) and color by blue all the other edges. ■



**Lemma 6.** (i) *Let  $t \geq 6$  and  $p \in \{2, 4, \dots, t\}$ . Then  $M_2(t) \in \mathcal{R}(K_{1,2}(v, v_p), C_4)$ .*

(ii) *Let  $t \geq 8$  and  $p \in \{4, 6, \dots, t - 4\}$ . Then  $M_2(t) \in \mathcal{R}(K_{1,2}(v_0, v_p), C_4)$ .*

The proof is similar to the previous proofs.

#### 4. MAIN RESULTS

Let  $n \geq 4$ . Let  $M_{a_j}, j = 1, 2, \dots, k$  be any graphs with roots  $r_{a_j,1}, r_{a_j,2}$  such that  $M_{a_j} \in \mathcal{R}(K_{1,2}(r_{a_j,1}, r_{a_j,2}), C_n)$ . Let  $L_{b_i}, i = 1, 2$  be any graphs with a root  $r_{b_i}$  such that  $L_{b_i} \in \mathcal{R}(K_{1,2}(r_{b_i}), C_n)$  and let  $L$  be any graph with roots  $r_1, r_2$ , where  $L \in \mathcal{R}(K_{1,2}(r_1, r_2)_1, C_n)$ .

Let  $P(a_1, a_2, \dots, a_k)$  be a graph which consists of  $k$  graphs  $M_{a_1}, M_{a_2}, \dots, M_{a_k}$ , where the vertex  $r_{a_j,2}$  is stuck to the vertex  $r_{a_{j+1},1}, j = 1, 2, \dots, k - 1$ . A graph  $C(a_1, a_2, \dots, a_k)$  is defined in the same way with the only difference that  $r_{a_1,1}$  is stuck to  $r_{a_k,2}$  as well.

Finally, we define the following families of graphs:

$B_1(C(a'_1, a'_2, \dots, a'_{k_1}), P(a_1, a_2, \dots, a_{k_2})), k_1 \geq n + 1, k_2 \geq 1$ , is a graph that consists of the graphs  $C(a'_1, a'_2, \dots, a'_{k_1})$  and  $P(a_1, a_2, \dots, a_{k_2})$ , where the first root of  $M_{a_1}$  is stuck to any root  $x$  of  $C(a'_1, a'_2, \dots, a'_{k_1})$  and the second root of  $M_{a_{k_2}}$  is stuck to any root  $y$  of  $C(a'_1, a'_2, \dots, a'_{k_1})$ , where  $d_{C(a'_1, a'_2, \dots, a'_{k_1})}(x, y) + d_{P(a_1, a_2, \dots, a_{k_2})}(x, y) \geq n + 1$ .

$B_2(L, P(a_1, a_2, \dots, a_k)), k \geq n$ , is a graph which consists of the graphs  $L$  and  $P(a_1, a_2, \dots, a_k)$ , where the first root of  $M_{a_1}$  is stuck to the first root of  $L$  and the second root of  $M_{a_k}$  is stuck to the second root of  $L$ .

$B_3(L_{b_1}, P(a_1, a_2, \dots, a_k), L_{b_2}), k \geq 0$ , is obtained by sticking the first root of  $M_{a_1}$  to the root of  $L_{b_1}$  and the second root of  $M_{a_k}$  is stuck to the root of  $L_{b_2}$ .

$B_4(C(a'_1, a'_2, \dots, a'_{k_1}), P(a_1, a_2, \dots, a_{k_2}), C(a''_1, a''_2, \dots, a''_{k_3})); k_1, k_3 \geq n + 1, k_2 \geq 0$ , is constructed by sticking the first root of  $M_{a_1}$  to any root of  $C(a'_1, a'_2, \dots, a'_{k_1})$  and the second root of  $M_{a_{k_2}}$  is stuck to any root of  $C(a''_1, a''_2, \dots, a''_{k_3})$ .

$B_5(L_{b_1}, P(a_1, a_2, \dots, a_{k_1}), C(a'_1, a'_2, \dots, a'_{k_2})), k_1 \geq 0, k_2 \geq n + 1$ , is obtained by sticking the first root of  $M_{a_1}$  to the root of  $L_{b_1}$  and the second root of  $M_{a_{k_1}}$  is stuck to any root of  $C(a'_1, a'_2, \dots, a'_{k_2})$ .

The graphs defined above will be also denoted briefly by  $B_1, B_2, \dots, B_5$ . The graphs  $M_{a'_i}, i = 1, 2, \dots, k_1$  and  $M_{a_j}, j = 1, 2, \dots, k_2$  will be called *seeds* of  $B_1$ . Seeds of  $B_2, B_3, B_4$  and  $B_5$  can be defined analogously. We show that  $B_1, B_2, \dots, B_5$  are Ramsey  $(K_{1,2}, C_n)$ -minimal graphs.

**Theorem 1.**  $B_1 \in \mathcal{R}(K_{1,2}, C_n)$ .

*Proof.* First let us show by contradiction that  $B_1 \rightarrow (K_{1,2}, C_n)$ . Assume that  $B_1 \not\rightarrow (K_{1,2}, C_n)$ . Since  $M_{a'_i} \in \mathcal{R}(K_{1,2}(r_{a'_i,1}, r_{a'_i,2}), C_n), i = 1, 2, \dots, k_1$  and  $M_{a_j} \in \mathcal{R}(K_{1,2}(r_{a_j,1}, r_{a_j,2}), C_n), j = 1, 2, \dots, k_2$ , by part (i) of Definition 2, we must color by red at least one edge incident to some root in  $M_{a'_i}$  (in  $M_{a_j}$ ) to have a red-blue coloring of the edges of  $M_{a'_i}$  (of  $M_{a_j}$ ) that contains neither a red copy of  $K_{1,2}$  nor a blue copy of  $C_n$ . But then, we have at least  $k_1 + k_2$  red edges incident to roots in  $B_1$ . Because the number of different roots in  $B_1$  is  $k_1 + k_2 - 1$ , there must be a red copy of  $K_{1,2}$  in any coloring of  $B_1$ . A contradiction.

In order to prove the minimality of  $B_1$  it suffices to show that  $B_1^* \not\rightarrow (K_{1,2}, C_n)$ , where  $B_1^* \simeq B_1 \setminus \{e\}$  for any fixed edge  $e \in E(B_1)$ . Suppose  $e \in E(M_{a'_i})$  where  $i \in \{1, 2, \dots, k_1\}$ . (The case  $e \in E(M_{a_j}), j \in \{1, 2, \dots, k_2\}$  can be handled similarly). Then  $M_{a'_i}^* \simeq M_{a'_i} \setminus \{e\}$ . We know that  $M_{a'_i}^* \not\rightarrow (K_{1,2}(r_{a'_i,1}, r_{a'_i,2}), C_n)$ , which means that there exists a red-blue coloring of the edges of  $M_{a'_i}^*$  containing neither a red copy of  $K_{1,2}$  nor a blue copy of  $C_n$  such that the roots  $r_{a'_i,1}, r_{a'_i,2}$  are not incident to red edges in  $M_{a'_i}^*$ .

From Definition 2 it follows that in any other seed of  $B_1^*$  we must color by red some edges incident to any fixed root, while the second root does not have to be incident to red edges of the seed to have a red-blue coloring of the seed containing no red  $K_{1,2}$  and no blue  $C_n$ . Note that since the coloring contains no red  $K_{1,2}$ , there must be just one red edge in the seed which is incident to the fixed root.

Thus, we can color the edges of  $B_1^*$  such that every root is incident to exactly one red edge. We do not have any red copy of  $K_{1,2}$  in the coloring of  $B_1^*$ . Since the number of seeds in  $C(a'_1, a'_2, \dots, a'_{k_1})$  is  $k_1 \geq n + 1$  and  $d_{C(a'_1, a'_2, \dots, a'_{k_1})}(x, y) + d_{P(a_1, a_2, \dots, a_{k_2})}(x, y) \geq n + 1$ , we do not have any blue copy of  $C_n$  in the coloring of  $B_1^*$  as well. This finishes the proof. ■

**Theorem 2.**  $B_2 \in \mathcal{R}(K_{1,2}, C_n)$ .

*Proof.* We show that  $B_2 \rightarrow (K_{1,2}, C_n)$ . Suppose the contrary, let  $B_2 \not\rightarrow (K_{1,2}, C_n)$ . Since  $M_{a_i} \in \mathcal{R}(K_{1,2}(r_{a_i,1}, r_{a_i,2}), C_n), i = 1, 2, \dots, k$  and  $L \in \mathcal{R}(K_{1,2}(r_1, r_2)_1, C_n)$ , from part (i) of Definition 2 it follows that we must have at least one red edge incident to some root in  $M_{a_i}$  to obtain a red-blue coloring of the edges of  $M_{a_i}$  containing neither a red copy of  $K_{1,2}$  nor a blue copy of  $C_n$ .

In any red-blue coloring of  $L$  that contains no red  $K_{1,2}$  and no blue  $C_n$ , there must be at least one red edge  $e_1$  incident to the first root in  $L$  and at least one red edge  $e_2$  incident to the second root in  $L$ , where the edges  $e_1, e_2$  are not necessarily different. Because the number of different roots in  $B_2$  is  $k + 1$ , there must be a root incident to at least two red edges. We have a red copy of  $K_{1,2}$  in the coloring of  $B_2$ , a contradiction.

Let us prove that  $B_2^* \not\rightarrow (K_{1,2}, C_n)$  for  $B_2^* \simeq B_2 \setminus \{e\}$ , where  $e$  is any fixed edge of  $B_2$ . We distinguish two cases:

a) Let  $e \in E(M_{a_i})$  where  $i \in \{1, 2, \dots, k\}$ . Then  $M_{a_i}^* \simeq M_{a_i} \setminus \{e\}$  and  $M_{a_i}^* \not\rightarrow (K_{1,2}(r_{a_i,1}, r_{a_i,2}), C_n)$ , which says that there exists a red-blue coloring of  $M_{a_i}^*$  containing neither a red  $K_{1,2}$  nor a blue  $C_n$ , where there are no red edges incident to the roots  $r_{a_i,1}, r_{a_i,2}$  in  $M_{a_i}^*$ .

Now consider all the other seeds  $M_{a_j}, j = 1, 2, \dots, k, j \neq i$  and  $L$ . By Definition 2, in any seed  $M_{a_j}$  we must color by red some edges incident to any fixed root to have a red-blue coloring of  $M_{a_j}$  that contains neither a red  $K_{1,2}$  nor a blue  $C_n$ . The second root does not have to be incident to any red edge of  $M_{a_j}$ . Since the coloring does not contain any red  $K_{1,2}$ , the fixed root is incident to exactly one red edge in  $M_{a_j}$ . In the seed  $L$ , if we have exactly one red edge incident to the first root and one red edge incident to the second root, there exists a red-blue coloring of  $L$  that does not contain any red  $K_{1,2}$  and any blue  $C_n$ .

It follows that it is possible to color the edges of  $B_2^*$  such that every root is incident to exactly one red edge, hence there is no red  $K_{1,2}$  in the coloring of  $B_2^*$ . Because the number of seeds in  $B_2^*$  is  $k + 1 \geq n + 1$ , there is also no blue  $C_n$  in the coloring.

b) Let  $e \in E(L)$ . Then  $L^* \simeq L \setminus \{e\}$  and  $L^* \not\rightarrow (K_{1,2}(r_j), C_n), j = 1, 2$ , which means that there is a red-blue coloring of  $L^*$  that contains neither a red  $K_{1,2}$  nor a blue  $C_n$ , where there is no red edge incident to  $r_j$  in  $L^*$ . Note that the other root can be incident to at most one red edge in  $L^*$ , otherwise we have a red  $K_{1,2}$  in the coloring of  $L^*$ .

Consider the seeds  $M_{a_j}, j = 1, 2, \dots, k$ . Analogously as in case a) it suffices to color by red exactly one edge of  $M_{a_j}$  which is incident to any root, while the second root does not have to be incident to any red edge in  $M_{a_j}$  to have a red-blue coloring of  $M_{a_j}$  that contains no red  $K_{1,2}$  and no blue  $C_n$ . Then we are able to color  $B_2^*$  such that we have neither a red  $K_{1,2}$  nor a blue  $C_n$  in the coloring. The proof is complete. ■

**Theorem 3.**  $B_5 \in \mathcal{R}(K_{1,2}, C_n)$ .

**Proof.** Let us prove by contradiction that  $B_5 \rightarrow (K_{1,2}, C_n)$ . Because  $M_{a_i} \in \mathcal{R}(K_{1,2}(r_{a_i,1}, r_{a_i,2}), C_n)$ ,  $i = 1, 2, \dots, k_1$  (because  $M_{a'_j} \in \mathcal{R}(K_{1,2}(r_{a'_j,1}, r_{a'_j,2}), C_n)$ ,  $j = 1, 2, \dots, k_2$  and  $L_{b_1} \in \mathcal{R}(K_{1,2}(r_{b_1}), C_n)$ ), in any red-blue coloring of  $M_{a_i}$  (of  $M_{a'_j}$ ,  $L_{b_1}$ ) that contains no red  $K_{1,2}$  and no blue  $C_n$ , there must be at least one red edge incident to some root in  $M_{a_i}$  (in  $M_{a'_j}$ ,  $L_{b_1}$ ). Then there are at least  $k_1 + k_2 + 1$  red edges incident to roots in  $B_5$ . Since the number of roots in  $B_5$  is  $k_1 + k_2$ , we have a red  $K_{1,2}$  in any coloring of  $B_5$ . A contradiction.

We show that  $B_5^* \not\rightarrow (K_{1,2}, C_n)$  for the graph  $B_5^* \simeq B_5 \setminus \{e\}$ , where  $e$  is any fixed edge of  $B_5$ . Assume that  $e \in E(M_{a_i})$  where  $i \in \{1, 2, \dots, k_1\}$ . (The cases  $e \in E(M_{a'_j})$ ,  $j \in \{1, 2, \dots, k_2\}$  and  $e \in E(L_{b_1})$  are similar.) Then  $M_{a_i}^* \simeq M_{a_i} \setminus \{e\}$  and  $M_{a_i}^* \not\rightarrow (K_{1,2}(r_{a_i,1}, r_{a_i,2}), C_n)$ , which means that there exists a red-blue coloring of  $M_{a_i}^*$  containing neither a red  $K_{1,2}$  nor a blue  $C_n$  such that  $r_{a_i,1}, r_{a_i,2}$  are not incident to red edges in  $M_{a_i}^*$ .

In any other seed of  $B_5^*$ , if one of the roots is not incident to red edges of the seed and the second root is incident to exactly one red edge, there exists a red-blue coloring of the seed that contains neither a red  $K_{1,2}$  nor a blue  $C_n$  (in  $L_{b_1}^*$  we have just one root which is incident to one red edge of  $L_{b_1}^*$ ).

Hence, it is possible to color the edges of  $B_5^*$  such that every root is incident to exactly one red edge and there is no red  $K_{1,2}$  in the coloring of  $B_5^*$ . Because the number of seeds in  $C(a'_1, a'_2, \dots, a'_{k_2})$  is  $k_2 \geq n + 1$ , there is no blue  $C_n$  in the coloring as well. ■

Similarly as Theorem 3, we can prove the next theorem.

**Theorem 4.**  $B_3, B_4 \in \mathcal{R}(K_{1,2}, C_n)$ .

Theorems 1–4 in combination with Lemmas 1–6 give infinite families of Ramsey  $(K_{1,2}, C_4)$ -minimal graphs.

For example, the graph  $B_3(L_m(t'_1), P(a_1, a_2, \dots, a_k), L_n(t'_2))$ , where  $P(a_1, a_2, \dots, a_k)$  consists of the graphs  $M_{a_j}(t_j)$ ,  $j = 1, 2, \dots, k$  and  $a_j, m, n \in \{1, 2, 3\}$  is a Ramsey  $(K_{1,2}, C_4)$ -minimal graph. Values of the parameters  $t'_1, t'_2, t_j$  follow from Lemmas 1–6.

Let  $B_3(L_m(t'_1), P(a_1, a_2, \dots, a_k), L_n(t'_2))$  contains exactly  $r$  seeds  $M_2(t_j)$ ,  $j \in \{1, 2, \dots, k\}$  such that the vertex which has degree  $t_j/2$  in  $M_2(t_j)$  is one of the roots of  $M_2(t_j)$  and let  $B_3(L_m(t'_1), P(a_1, a_2, \dots, a_k), L_n(t'_2))$  also contains  $z$  seeds  $L_3(6)$  with the root denoted by  $v_4$  in  $L_3(6)$ . Note

that  $0 \leq r \leq k$  and  $0 \leq z \leq 2$ . It is easy to show that the diameter of  $B_3(L_m(t'_1), P(a_1, a_2, \dots, a_k), L_n(t'_2))$  is  $2k + 6 - r - z$ , since

- the eccentricity of the root of  $L_i(t')$  is 3 for  $i = 1, 2, 3$  and any  $t'$  except for the eccentricity of  $v_4$  in  $L_3(6)$  that is equal to 2,
- the distance between two roots in  $M_i(t)$  is 2 for  $i = 1, 3$ , while in  $M_2(t)$  the roots can be adjacent.

It follows that we found an infinite class of Ramsey  $(K_{1,2}, C_4)$ -minimal graphs for every diameter  $\geq 4$ . The problem of existence of an infinite family of Ramsey  $(K_{1,2}, C_4)$ -minimal graphs of diameter 3 remains open.

### Acknowledgement

Research of the first author was supported by the VEGA Grant No. 1/2004/05 and the APVV Grants No. 20–000704 and 40–06.

### REFERENCES

- [1] E.T. Baskoro, L. Yulianti and H. Assiyatun, *Ramsey  $(K_{1,2}, C_4)$ -minimal graphs*, J. Combin. Mathematics and Combin. Computing **65** (2008) 79–90.
- [2] M. Borowiecki, M. Hałuszczak and E. Sidorowicz, *On Ramsey-minimal graphs*, Discrete Math. **286** (2004) 37–43.
- [3] M. Borowiecki, I. Schiermeyer and E. Sidorowicz, *Ramsey  $(K_{1,2}, K_3)$ -minimal graphs*, Electronic J. Combinatorics **12** (2005) #R20.
- [4] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, *Ramsey-minimal graphs for star-forests*, Discrete Math. **33** (1981) 227–237.
- [5] S.A. Burr, P. Erdős and L. Lovász, *On graphs of Ramsey type*, Ars Combin. **1** (1976) 167–190.
- [6] T. Łuczak, *On Ramsey-minimal graphs*, Electronic J. Combinatorics **1** (1994) #R4.
- [7] I. Mengersen and J. Oeckermann, *Matching-star Ramsey sets*, Discrete Appl. Math. **95** (1999) 417–424.

Received 3 March 2009

Revised 13 January 2010

Accepted 13 January 2010