

ON VERTEX STABILITY WITH REGARD TO COMPLETE BIPARTITE SUBGRAPHS

ANETA DUDEK AND ANDRZEJ ŻAK

Faculty of Applied Mathematics
AGH University of Science and Technology
Mickiewicza 30, 30-059 Kraków, Poland
e-mail: {dudekane,zakandr}@agh.edu.pl

Abstract

A graph G is called $(H; k)$ -vertex stable if G contains a subgraph isomorphic to H ever after removing any of its k vertices. $Q(H; k)$ denotes the minimum size among the sizes of all $(H; k)$ -vertex stable graphs. In this paper we complete the characterization of $(K_{m,n}; 1)$ -vertex stable graphs with minimum size. Namely, we prove that for $m \geq 2$ and $n \geq m + 2$, $Q(K_{m,n}; 1) = mn + m + n$ and $K_{m,n} * K_1$ as well as $K_{m+1,n+1} - e$ are the only $(K_{m,n}; 1)$ -vertex stable graphs with minimum size, confirming the conjecture of Dudek and Zwonek.

Keywords: vertex stable, bipartite graph, minimal size.

2010 Mathematics Subject Classification: 05C70, 11B50, 05C78.

1. INTRODUCTION

We deal with simple graphs without loops and multiple edges. We use the standard notation of graph theory, cf. [1]. The following notion was introduced in [2]. Let H be any graph and k a non-negative integer. A graph G is called $(H; k)$ -vertex stable if G contains a subgraph isomorphic to H ever after removing any of its k vertices. Then $Q(H; k)$ denotes minimum size among the sizes of all $(H; k)$ -vertex stable graphs. Note that if H does not have isolated vertices then after adding to or removing from a $(H; k)$ -vertex

The research of the authors was partially supported by the Polish Ministry of Science and Higher Education.

stable graph any number of isolated vertices we still have a $(H; k)$ -vertex stable graph with the same size. Therefore, in the sequel we assume that no graph in question has isolated vertices.

There are two trivial examples of (H, k) -vertex stable graphs, namely $(k+1)H$ (a disjoint union of $(k+1)$ copies of H) and $H * K_k$ (a graph obtained from $H \cup K_k$ by joining all the vertices of H to all the vertices of K_k). Therefore,

Proposition 1. $Q(H; k) \leq \min \left\{ (k+1)|E(H)|, |E(H)| + k|V(H)| + \binom{k}{2} \right\}.$

On the other hand, the following is easily seen.

Proposition 2. *Suppose that H contains k vertices which cover q edges. Then $Q(H; k) \geq |E(H)| + q$.*

Recall also the following

Proposition 3 ([2]). *Let δ_H be a minimal degree of a graph H . Then in any $(H; k)$ -vertex stable graph G with minimum size, $\deg_G v \geq \delta_H$ for each vertex $v \in G$.*

The exact values of $Q(H; k)$ are known in the following cases: $Q(C_i; k) = i(k+1)$, $i = 3, 4$, $Q(K_4; k) = 5(k+1)$, $Q(K_n; k) = \binom{n+k}{2}$ for n large enough, and $Q(K_{1,m}; k) = m(k+1)$, $Q(K_{n,n}; 1) = n^2 + 2n$, $Q(K_{n,n+1}; 1) = (n+1)^2$, $n \geq 2$, see [2, 3]. In this paper we complete the characterization of $(K_{m,n}; 1)$ vertex stable graphs with minimum size. Namely, we prove the following theorem and hence confirm Conjecture 1 formulated in [3].

Theorem 1. *Let m, n be positive integers such that $m \geq 2$ and $n \geq m+2$. Then $Q(K_{m,n}; 1) = mn + m + n$ and $K_{m,n} * K_1$ as well as $K_{m+1,n+1} - e$, where $e \in E(K_{m+1,n+1})$, are the only $(K_{m,n}; 1)$ -vertex stable graphs with minimum size.*

2. PROOF OF THE MAIN RESULT

Proof of Theorem 1. Let $m \geq 2$ and $n \geq m+2$ be positive integers. Define $G_1 := K_{m,n} * K_1$ and $G_2 := K_{m+1,n+1} - e$ where $e \in E(K_{m+1,n+1})$. Let $G = (V, E)$ be a $(K_{m,n}, 1)$ -vertex stable graph with minimum size. Thus, by Proposition 1, $|E(G)| \leq mn + m + n$. Clearly G contains a subgraph

H isomorphic to $K_{m,n}$. Let $H = (X, Y; E_H)$ with vertex bipartition sets X, Y such that $|X| = m$ and $|Y| = n$. Let $v \in X$. Since G is $(K_{m,n}; 1)$ -vertex stable, $G - v$ contains a subgraph H' isomorphic to $K_{m,n}$. Let $H' = (X', Y'; E_{H'})$ with vertex bipartition sets X', Y' such that $|X'| = m$ and $|Y'| = n$. We denote $x_1 = |X \cap X'|$, $x_2 = |X \cap Y'|$, $y_1 = |Y \cap X'|$, $y_2 = |Y \cap Y'|$. Hence $x_1 + x_2 \leq m - 1$, $y_1 + y_2 \leq n$, $y_1 \leq m$. One can see that $|E(G)| \geq 2mn - x_1y_2 - x_2y_1$. Consider the following linear programming problem with respect to y_1 and y_2

$$\begin{cases} y_1 \leq m \\ y_1 + y_2 \leq n \\ y_1 \geq 0 \\ y_2 \geq 0 \\ c = x_1y_2 + x_2y_1 \rightarrow \max \end{cases}$$

where x_1 and x_2 are parameters such that $x_1, x_2 \geq 0, x_1 + x_2 \leq m - 1$.




Figure 1. Geometrical interpretation of the linear programming problem.

The proof falls into two cases.

Case 1. $x_1 < x_2$.

In this case $y_1 = m, y_2 = n - m, c = x_2m + x_1(n - m)$ is the unique optimal solution of the above linear programming problem. This can be easily checked using a geometrical interpretation of the linear programming problem, see Figure 1. Thus $|E(G)| \geq 2mn - x_2m - x_1(n - m)$ and the inequality is strict if $y_1 \neq m$ or $y_2 \neq n - m$. We assume that $x_1 + x_2 = m - 1$ because otherwise the size of G may only increase. Then

$$|E(G)| \geq 2mn - m^2 + m + x_1(2m - n) := f(x_1).$$

Subcase 1a. $n > 2m$.

Then $f(x_1)$ is decreasing. Furthermore, $x_1 < \frac{m-1}{2}$ since $x_1 < x_2$. Thus

$$|E(G)| > f\left(\frac{m-1}{2}\right) = \frac{3}{2}mn + \frac{1}{2}n \geq mn + m + n.$$

Thus $|E(G)| > mn + m + n$, a contradiction.

Subcase 1b. $n < 2m$.

Then $f(x_1)$ is increasing. Thus

$$E(G) \geq f(0) = 2mn - m^2 + m \geq mn + m + n$$

with equality if and only if $m = 2$ and $n = 4$, which is not possible in this subcase.

Subcase 1c. $n = 2m$.

In this case

$$E(G) \geq mn + m + n$$

with equality if and only if $m = 2, n = 4, y_1 = y_2 = 2$. Recall that $x_1 < x_2$ whence $x_1 = 0$ and $x_2 = 1$. Let $u \in Y' \setminus (X \cup Y)$. Thus $|E(G)| \geq 12 + \deg u$. Hence $\deg u = 2$ and $|V(G)| = 7$ because otherwise $|E(G)| > mn + m + n$. However, then G is not $(K_{2,4}; 1)$ -stable. Indeed let w be a neighbor of u . Then $G - w$ does not contain any subgraph isomorphic to $K_{2,4}$ since $G - w$ has 6 vertices and one of them has degree 1. Therefore Case 1 is not possible.

Case 2. $x_1 \geq x_2$.

In this case $c = x_1n$ is the optimal solution of the above linear problem, see Figure 1. Therefore, $|E(G)| \geq 2mn - x_1n$. If $x_1 \leq m - 2$ then $|E(G)| \geq 2mn - (m - 2)n = mn + 2n > mn + m + n$. Hence we may assume that

$x_1 = m - 1$ and $x_2 = 0$. Thus there is only one vertex, say u , such that $u \in X' \setminus X$.

Subcase 2a. $y_2 = n$.

Thus, u have n neighbors in Y . Note that $|V(G)| \leq m + n + 2$. Indeed, otherwise by Proposition 3, $|E(G)| \geq mn + n + 2m - 1 > mn + m + n$. Consider now a graph $G'' := G - w$ where $w \in Y$. Clearly $G - w$ contains a subgraph H'' isomorphic to $K_{m,n}$. Let $H'' = (X'', Y''; E_{H''})$ with vertex bipartition sets X'', Y'' such that $|X''| = m$ and $|Y''| = n$. Let $x'_1 = |X \cap X''|$, $x'_2 = |X \cap Y''|$, $y'_1 = |Y \cap X''|$, $y'_2 = |Y \cap Y''|$.

Suppose first that $|V(G)| = m + n + 2$ and $u, u_1 \in V(G) \setminus (X \cup Y)$. Since $|E(G)| \leq mn + m + n$, $\deg u_1 = m$ and $\deg u \leq n + 1$. In particular, $u_1 \notin X''$ and u has no neighbor in X . Furthermore, $|E(G)| \geq mn + n + m + x'_1 x'_2 + y'_1 y'_2$. Thus, $x'_1 = 0$ or $x'_2 = 0$, and $y'_1 = 0$ or $y'_2 = 0$. We distinguish two possibilities

1. $x'_1 = 0$. Then $y'_1 \neq 0$. Indeed, otherwise $X'' = \{u, u_1\}$, a contradiction with previous observation that $u_1 \notin X''$. Hence, $y'_2 = 0$. Thus, $x'_2 = m$ and $u, u_1 \in Y''$ (so $n = m + 2$). Consequently, $y'_1 = m$. However, then G is not $(K_{m,m+2}; 1)$ -stable. Indeed, let w_1 be a neighbor of u_1 , $w_1 \in X'' \subset Y$. Then $G - w_1$ consists of a subgraph isomorphic to $K_{m+1,m+1}$ plus one vertex (namely u_1) and $m - 1$ edges incident to it. Therefore, $G - w_1$ does not contain any subgraph isomorphic to $K_{m,m+2}$.

2. $x'_1 \neq 0$. Then $x'_2 = 0$ and $u \notin Y''$. Consequently, $u_1 \in Y''$ and $y'_2 \neq 0$. Hence $y'_1 = 0$. It is easy to see now that $G \cong G_2$.

Assume now that $|V(G)| = m + n + 1$. Hence $x'_1 + x'_2 = m$ and $y'_1 + y'_2 = n - 1$. We have the next two possibilities.

3. $x'_1 + y'_1 = m$. Then $|E(G)| \geq mn + x'_1 x'_2 + y'_1 y'_2 + \deg u \geq mn + x'_1 x'_2 + y'_1 y'_2 + n + x'_1$. Hence

$$|E(G)| \geq mn + (m - x'_1)(n - 1 - m + 2x'_1) + n + x'_1 =: f_1(x'_1), \quad 0 \leq x'_1 \leq m.$$

It is not difficult to see that $f_1(x'_1)$ obtains the smallest value for $x'_1 = 0$ or $x'_1 = m$ only. Thus, $|E(G)| \geq \min\{f_1(0), f_1(m)\}$. Note that $f_1(0) = 2mn + n - m - m^2 \geq mn + m + n$ with equality if and only if $n = m + 2$. However, then there is a vertex $y \in Y''$ such that $G - y \cong K_{m+1,m+1}$ so $G - y$ does not contain any subgraph isomorphic to $K_{m,m+2}$. Furthermore, $f_1(m) \geq mn + n + m$. Thus, $|E(G)| \geq mn + m + n$ with equality if and only if $x_1 = m$. Then $G \cong G_1$.

4. $x'_2 + y'_2 = n$. Then $|E(G)| \geq mn + x'_1x'_2 + y'_1y'_2 + n + x'_2$. Hence,

$$|E(G)| \geq mn + (m - x'_2)x'_2 + (x'_2 - 1)(n - x'_2) + n + x'_2 =: f_2(x_2), \quad 1 \leq x'_2 \leq m.$$

One can see that $f_2(x'_2)$ obtains the smallest value for $x'_2 = 1$ or $x'_2 = m$ only. Thus, $|E(G)| \geq \min\{f_2(1), f_2(m)\}$. Note that $f_2(1) = mn + n + m$. Then $G \cong G_1$. On the other hand, $f_2(m) = 2mn + 2m - m^2 > mn + m + n$.

Subcase 2b. $y_2 < n$.

Thus, there is a vertex $z \in Y'$ such that $z \in V(G) \setminus (X \cup Y)$. This clearly forces $m - 1$ neighbours of z in $X \setminus \{v\}$. Consider now a graph $G - v_1$, $v \neq v_1 \in X$. We repeat all preceding arguments to the graph $G - v_1$. Consequently, $G \cong G_i$, $i = 1, 2$, or there is a vertex $z_1 \in V(G) \setminus (X \cup Y)$ which has $m - 1$ neighbors in $X \setminus \{v_1\}$. If $z = z_1$ then z has m neighbors in X and $G \cong G_1$ if $u \in Y$ or $G \cong G_2$ otherwise. If $z \neq z_1$ then either $\deg z + \deg z_1 \geq 2m + 1$ if both vertices z and z_1 are involved in a $K_{m,n}$ contained in $G - v$ or $G - v_1$, or $\deg u \geq n + 1$ otherwise. Thus, $|E(G)| \geq mn + 2m - 1 + n > mn + m + n$. ■

3. CONCLUDING REMARKS

In [2, 3] it is proved that $Q(K_{1,n}; k) = (k + 1)n$. However, for $n \geq 3$ the extremal graphs are not characterized.

Proposition 4. *Let G be a $(K_{1,n}; k)$ -vertex stable graph with minimum size, $n \geq 3$. Then $G = (k + 1)K_{1,n}$.*

Proof. The proof is by induction on k . The thesis is obvious for $k = 0$. Assume that $k > 0$. Let G be a $(K_{1,n}; k)$ -vertex stable graph with minimum size. Hence, $|E(G)| = (k + 1)n$. Note that each $(K_{1,n}; k)$ -vertex stable graph contains $k + 1$ vertices of degree at least n . Let $v \in V(G)$ with $\deg v \geq n$. Thus, $G - v$ is $(K_{1,n}; k - 1)$ -vertex stable graph with $|E(G - v)| \leq kn$. Hence, $|E(G - v)| = kn$ and $\deg v = n$. By the induction hypothesis $G - v = kK_{1,n}$. Note that v is not a neighbor of any vertex of degree n . Suppose on the contrary, that $uv \in E(G)$ and $\deg u = n$. Then $G - u$ contains only $k - 1$ vertices of degree greater than or equal to n whence is not $(K_{1,n}; k - 1)$ -vertex stable, a contradiction. Thus, G contains $k + 1$ independent vertices of degree exactly n . We will show that these vertices have pairwise disjoint sets of neighbors. Indeed, otherwise let x be a common neighbor of two

vertices of degree n . Thus, again, $G - x$ has only $k - 1$ vertices of degree greater than or equal to n , a contradiction. ■

In the following table we present the complete characterization of $(K_{m,n}; 1)$ -vertex stable graphs with minimum size.

$m; n$	$Q(K_{m,n}; 1)$	All $(K_{m,n}; 1)$ -vertex stable graphs with minimum size
$m = 1, n = 1$	2 [3]	$2K_{1,1}$ [3]
$m = 1, n = 2$	4 [3]	$K_{2,2}, 2K_{1,2}$ [3]
$m = 1, n \geq 3$	$2n$ [2]	$2K_{1,n}$
$m = 2, n = 2$	8 [3]	$K_{2,2} * K_1, K_{3,3} - e, 2K_{2,2}$ [3]
$m \geq 2, n = m + 1$	$(m + 1)^2$ [3]	$K_{m+1, m+1}$ [3]
$m \geq 3, n = m$	$m^2 + 2m$ [3]	$K_{m,m} * K_1, K_{m+1, m+1} - e$ [3]
$m \geq 2, n \geq m + 2$	$mn + m + n$	$K_{m,n} * K_1, K_{m+1, n+1} - e$

Acknowledgement

The research was partially supported by the University of Science and Technology grant No. 11 420 04. Authors wish to express his gratitude to Gyula Y. Katona for suggesting the problem.

REFERENCES

- [1] R. Diestel, Graph Theory, second ed. (Springer-Verlag, 2000).
- [2] A. Dudek, A. Szymański and M. Zwonek, (H, k) stable graphs with minimum size, Discuss. Math. Graph Theory **28** (2008) 137–149.
- [3] A. Dudek and M. Zwonek, (H, k) stable bipartite graphs with minimum size, Discuss. Math. Graph Theory **29** (2009) 573–581.

Received 23 October 2009

Revised 23 February 2010

Accepted 21 March 2010