

## PAIRED DOMINATION IN PRISMS OF GRAPHS

CHRISTINA M. MYNHARDT\*

AND

MARK SCHURCH

*Department of Mathematics and Statistics*  
*University of Victoria*  
*P.O. Box 3060 STN CSC*  
*Victoria, BC, Canada V8W 3R4*

**e-mail:** mynhardt@math.uvic.ca, mschurch@math.uvic.ca

### Abstract

The paired domination number  $\gamma_{\text{pr}}(G)$  of a graph  $G$  is the smallest cardinality of a dominating set  $S$  of  $G$  such that  $\langle S \rangle$  has a perfect matching. The generalized prisms  $\pi G$  of  $G$  are the graphs obtained by joining the vertices of two disjoint copies of  $G$  by  $|V(G)|$  independent edges. We provide characterizations of the following three classes of graphs:  $\gamma_{\text{pr}}(\pi G) = 2\gamma_{\text{pr}}(G)$  for all  $\pi G$ ;  $\gamma_{\text{pr}}(K_2 \square G) = 2\gamma_{\text{pr}}(G)$ ;  $\gamma_{\text{pr}}(K_2 \square G) = \gamma_{\text{pr}}(G)$ .

**Keywords:** domination, paired domination, prism of a graph, Cartesian product.

**2010 Mathematics Subject Classification:** 05C69.

### 1. INTRODUCTION

The *paired domination number* of a graph  $G$  is the smallest cardinality of a dominating set  $S$  of  $G$  such that  $\langle S \rangle$  has a perfect matching, and is denoted by  $\gamma_{\text{pr}}(G)$ . The paired domination number of the Cartesian product  $G \square H$  of two isolate-free graphs  $G$  and  $H$  was first investigated by Brešar, Henning and Rall [1], who obtained upper bounds on  $\gamma_{\text{pr}}(G)\gamma_{\text{pr}}(H)$  in terms

---

\*Supported by the Natural Sciences and Engineering Research Council of Canada.

of  $\gamma_{\text{pr}}(G \square H)$ . They showed, i.e., that for any nontrivial tree  $T$  and any isolate-free graph  $H$ ,  $\gamma_{\text{pr}}(T)\gamma_{\text{pr}}(H) \leq 2\gamma_{\text{pr}}(T \square H)$ .

We compare the paired domination number of a graph  $G$  with the paired domination numbers of its generalized prisms  $\pi G$ ; i.e., the graphs obtained by joining the vertices of two disjoint copies of  $G$  by  $|V(G)|$  independent edges. Obviously,  $\gamma_{\text{pr}}(\pi G) \leq 2\gamma_{\text{pr}}(G)$ . Graphs  $G$  for which  $\gamma_{\text{pr}}(\pi G) = 2\gamma_{\text{pr}}(G)$  regardless of how the two copies of  $G$  are joined are called *universal  $\gamma_{\text{pr}}$ -doubblers*.

After providing background information in Section 2, we give necessary and sufficient conditions for a graph to be a universal  $\gamma_{\text{pr}}$ -doubler in Section 3. We also give necessary and sufficient conditions for a graph to be a *prism  $\gamma_{\text{pr}}$ -doubler*, i.e.,  $\gamma_{\text{pr}}(K_2 \square G) = 2\gamma_{\text{pr}}(G)$  (Section 4), and a *prism  $\gamma_{\text{pr}}$ -fixer*, i.e.,  $\gamma_{\text{pr}}(K_2 \square G) = \gamma_{\text{pr}}(G)$  (Section 5). Open problems related to this topic are mentioned in Section 6.

## 2. DEFINITIONS AND BACKGROUND

For any permutation  $\pi$  of  $V(G)$ , the *prism of  $G$  with respect to  $\pi$*  is the graph  $\pi G$  obtained from two copies  $G_1$  and  $G_2$  of  $G$  by joining  $u \in V(G_1)$  and  $v \in V(G_2)$  if and only if  $v = \pi(u)$ . If  $\pi$  is the identity  $\mathbf{1}_G$ , then  $\pi G = K_2 \square G$ , the *Cartesian product* of  $G$  and  $K_2$ . The graph  $K_2 \square G$  is called the *prism of* (or *over*)  $G$  and, in general,  $\pi G$  is a *generalized prism* of  $G$ .

We shall abbreviate  $V(G)$ ,  $E(G)$  and  $V(G_i)$  to  $V$ ,  $E$  and  $V_i$ , respectively. Let  $u \in V$  and  $S \subseteq V$ . In  $\pi G$  we denote the counterparts of  $u$  (or  $S$ ) in  $G_1$  and  $G_2$  by  $u_1$  and  $u_2$  (or  $S_1$  and  $S_2$ ) respectively. Conversely, the vertex  $u_1$  and set  $S_1$  in  $G_1$  (or  $u_2$  and  $S_2$  in  $G_2$ ) are denoted by  $u$  and  $S$  respectively when considered in  $G$ .

For  $v \in V$ , the *open neighbourhood*  $N(v)$  of  $v$  is defined by  $N(v) = \{u \in V : uv \in E\}$ , and the *closed neighbourhood*  $N[v]$  of  $v$  is the set  $N(v) \cup \{v\}$ . For  $S \subseteq V$ ,  $N(S) = \bigcup_{s \in S} N(s)$ ,  $N[S] = \bigcup_{s \in S} N[s]$  and  $N\{S\} = N[S] - S$ . For  $v \in S$  we call  $w \in V - S$  an  *$S$ -external private neighbour* of  $v$  if  $N(w) \cap S = \{v\}$ . Denote the set of all  $S$ -external private neighbours of  $v$  by  $\text{epn}(v, S)$ .

A set  $S \subseteq V$  *dominates*  $G$  or is a *dominating set* of  $G$  if every vertex in  $V - S$  is adjacent to a vertex in  $S$ . The *domination number*  $\gamma(G)$  of  $G$  is defined by  $\gamma(G) = \min\{|S| : S \text{ dominates } G\}$ . A dominating set  $S$  is a *paired dominating set (PDS)* if  $\langle S \rangle$  has a perfect matching. A vertex  $v$  is an  *$\overline{M}$ -vertex* of a matching  $M$  if  $v$  does not belong to any edge of  $M$ . If  $S$

is a PDS and  $M$  is a perfect matching of  $\langle S \rangle$ , we call  $M$  an  $S$ -matching. A  $\gamma$ -set of  $G$  is a dominating set of  $G$  of cardinality  $\gamma(G)$ ; a  $\gamma_{\text{pr}}$ -set is defined similarly. We follow [9] for domination terminology.

It is easy to see that  $\gamma(G) \leq \gamma(\pi G) \leq 2\gamma(G)$  for all permutations  $\pi$  of  $V$ . If  $\gamma(K_2 \square G) = \gamma(G)$ , then  $G$  is called a *prism fixer*, and if  $\gamma(K_2 \square G) = 2\gamma(G)$ , then  $G$  is a *prism doubler*. If  $\gamma(\pi G) = \gamma(G)$  for all permutations  $\pi$  of  $V$ , then  $G$  is a *universal fixer*, and if  $\gamma(\pi G) = 2\gamma(G)$  for all  $\pi$ , then  $G$  is a *universal doubler*.

Prism fixers we first studied by Hartnell and Rall [7, 8] in connection with Vizing's conjecture on the domination number of the Cartesian product of graphs. Prism and universal doublers were studied in [3], while fixers and doublers for other domination parameters, such as total and paired domination, were investigated in [11]. The graphs  $\overline{K_n}$ ,  $n \geq 1$ , are universal fixers because  $\pi \overline{K_n} = nK_2$  for all permutations  $\pi$  of  $V$ . Moreover, these graphs are the only universal fixers known to date. The following conjecture was formulated in [10] and also studied in [2, 4, 6].

**Conjecture 1.** *The graphs  $\overline{K_n}$ ,  $n \geq 1$ , are the only universal fixers.*

It is obvious that  $\gamma_{\text{pr}}(\pi G) \leq 2\gamma_{\text{pr}}(G)$  for any graph  $G$  and any permutation  $\pi$  of  $V$ . Unlike the case for the domination number, though, the paired domination number of  $\pi G$  is not bounded below by the paired domination number of  $G$ . For the graph  $G$  in Figure 1,  $\gamma_{\text{pr}}(G) = 6$ , but for any  $\pi G$  obtained by adding enough edges to the graph shown,  $\gamma_{\text{pr}}(\pi G) = 4$ . However, if  $\pi$  is the identity, then the above-mentioned lower bound follows from the work in [1]. We give a direct proof below.




Figure 1.  $\gamma_{\text{pr}}(\pi G) < \gamma_{\text{pr}}(G)$

**Proposition 1.** *For any isolate-free graph  $G$ ,  $\gamma_{\text{pr}}(G) \leq \gamma_{\text{pr}}(K_2 \square G) \leq 2\gamma(G)$ .*

**Proof.** For the upper bound, note that if  $D$  is a  $\gamma$ -set of  $G$ , then  $D_1 \cup D_2$  is a PDS of  $K_2 \square G$ . For the lower bound, let  $W$  be a  $\gamma_{\text{pr}}$ -set of  $K_2 \square G$  with  $X_1 = W \cap V_1$  and  $D_2 = W \cap V_2$  and let  $S = X \cup D$ . Then  $S$  dominates  $G$  and  $|S| = |X| + |D| - |X \cap D|$ .

If  $X \cap D = \emptyset$ , then  $\langle S \rangle$  contains a perfect matching (the matching corresponding to the perfect matching of  $\langle W \rangle$ ) and  $S$  is a PDS of  $G$  with  $|S| = |W|$ , so we are done.

Assume  $X \cap D \neq \emptyset$ . Let  $M$  be a maximum matching of  $\langle S \rangle$  and  $Z = \{z^1, \dots, z^k\}$  the set of  $\overline{M}$ -vertices; note that  $k \leq |X \cap D|$ . Let  $S^0 = S$  and for  $i = 1, \dots, k$ , construct  $S^i$  recursively as follows.

- If  $z^i$  is adjacent to  $s^i \in V - S^{i-1}$ , let  $S^i = S^{i-1} \cup \{s^i\}$ . Otherwise,  $z^i$  is adjacent to  $x \in S^{i-1}$  because  $G$  is isolate-free; hence  $N[z^i] \subseteq S^{i-1}$ . Let  $S^i = S^{i-1} - \{z^i\}$ .

Then  $S^k$  dominates  $G$ ,  $\langle S^k \rangle$  has a perfect matching and thus  $S^k$  is a PDS of  $G$ . Moreover,  $|S^k| \leq |S| + |Z| \leq |X| + |D| = |W|$  and the result follows. ■

**Corollary 2.** *If a graph  $G$  is a prism  $\gamma_{\text{pr}}$ -doubler, then  $\gamma_{\text{pr}}(G) = \gamma(G)$ .*

### 3. UNIVERSAL DOUBLERS

Suppose  $D'$  is a  $\gamma_{\text{pr}}$ -set of a graph  $G$  in which  $u$  is paired with  $v$ , and  $\text{epn}(v, D') = \emptyset$ . Then  $D = D' - \{v\}$  dominates  $G$ , and  $D_1 \cup D_2$  is a  $\gamma_{\text{pr}}$ -set of  $K_2 \square G$  in which  $u_1$  is paired with  $u_2$ . Thus  $G$  is not a prism  $\gamma_{\text{pr}}$ -doubler and thus not a universal  $\gamma_{\text{pr}}$ -doubler. A similar argument (but with another permutation) shows that if  $G$  has a  $\gamma_{\text{pr}}$ -set  $D$  in which  $|\text{epn}(v, D)|$  is small compared to  $\gamma_{\text{pr}}(G)$  for some vertex  $v \in D$ , then  $G$  is not a universal  $\gamma_{\text{pr}}$ -doubler. These cases suggest that vertices contained in  $\gamma_{\text{pr}}$ -sets of universal  $\gamma_{\text{pr}}$ -doubblers have large degrees relative to  $\gamma_{\text{pr}}(G)$ , and hence that  $\gamma_{\text{pr}}(G)$  is small compared to the order of  $G$ , which we denote throughout by  $n$ .

In this section we obtain necessary and sufficient conditions for a graph to be a universal  $\gamma_{\text{pr}}$ -doubler. These conditions easily lead to an upper bound on the paired domination number of a universal  $\gamma_{\text{pr}}$ -doubler  $G$ , and lower bounds on the degrees and number of external private neighbours of the vertices in  $\gamma_{\text{pr}}$ -sets of  $G$ .

We begin with a simple lemma.

**Lemma 3.** *If  $\gamma(G) = \gamma_{\text{pr}}(G)$ , then  $n \geq 2\gamma_{\text{pr}}(G)$  and  $G$  has a PDS of cardinality  $\gamma_{\text{pr}}(G) + 2i$  for each  $1 \leq i \leq \gamma_{\text{pr}}(G)/2$ .*

**Proof.** It is well known [9, Theorem 2.1] that  $n \geq 2\gamma(G)$ , so  $n \geq 2\gamma_{\text{pr}}(G)$ . The latter part of the statement follows because each pair of vertices in a  $\gamma_{\text{pr}}$ -set  $X$  which is also a  $\gamma$ -set can be split into two pairs since each vertex of  $X$  has an external private neighbour [9, Theorem 1.1]. ■

We next define notation that will be used throughout this section. Let

$$(1) \quad \begin{cases} X \subseteq V \text{ such that } 0 < |X| < \gamma_{\text{pr}}(G); \\ Y = V - N[X]; \\ M \text{ be a matching of } \langle X \rangle; \\ Z = X - V(M), \text{ i.e., } Z \text{ is the set of } \overline{M}\text{-vertices in } X; \\ k = |Z|. \end{cases}$$

We now characterize universal  $\gamma_{\text{pr}}$ -doublers in terms of the cardinalities of the sets  $X$ ,  $Y$  and  $Z$  as defined in (1).

**Theorem 4.** *A graph  $G$  is a universal  $\gamma_{\text{pr}}$ -doubler if and only if, for each set  $X \subseteq V$  with  $0 < |X| < \gamma_{\text{pr}}(G)$ , a maximum matching  $M$  of  $\langle X \rangle$ , and  $Y$  and  $k$  as defined in (1),*

$$|Y| \geq 2\gamma_{\text{pr}}(G) - |X| - k - 1.$$

**Proof.** Suppose that for some  $X \subseteq V$  with  $0 < |X| < \gamma_{\text{pr}}(G)$ ,

$$|Y| < 2\gamma_{\text{pr}}(G) - |X| - k - 1.$$

We consider two cases, depending on the parity of  $k$ .

*Case 1.*  $k$  is even.

Then by definition of  $Z$ ,  $|X|$  is even. Choose a PDS  $D$  of  $G$  as follows.

- (i) If  $|Y| + k \leq \gamma_{\text{pr}}(G)$ , then let  $D$  be any  $\gamma_{\text{pr}}$ -set of  $G$ .
- (ii) Otherwise, let  $D$  be any PDS of  $G$  with  $|D| = |Y| + k$  if  $|Y|$  is even, or  $|D| = |Y| + k + 1$  if  $|Y|$  is odd. (A PDS of this size exists by Lemma 3.)

Let  $\pi$  be any permutation of  $V$  such that  $\pi(Y \cup Z) \subseteq D$  and  $\langle \pi(Z) \rangle$  has a perfect matching  $M'$  that is contained in a  $D$ -matching. Then  $W = X_1 \cup D_2$  dominates  $\pi G$  and  $\langle W \rangle$  has a  $W$ -matching in which each edge  $u_2 v_2$  in  $M'_2$  is replaced by two edges  $z_1 u_2$  and  $z'_1 v_2$ , where  $z, z' \in Z$ . (See Figure 2.)




Figure 2.  $X_1 \cup D_2$  dominates  $\pi G$ .

Therefore  $W$  is a PDS of  $\pi G$ . If  $D$  is a  $\gamma_{\text{pr}}$ -set of  $G$  (i.e., if  $D$  was defined in (i)), then

$$|W| = |X| + |D| < 2\gamma_{\text{pr}}(G),$$

i.e.,  $G$  is not a universal  $\gamma_{\text{pr}}$ -doubler. If  $D$  was defined in (ii), then

$$\begin{aligned} |W| &= |X| + |D| \\ &\leq |X| + |Y| + k + 1 \\ &< |X| + (2\gamma_{\text{pr}}(G) - |X| - k - 1) + k + 1 \\ &= 2\gamma_{\text{pr}}(G), \end{aligned}$$

and  $G$  is not a universal doubler in this case either.

*Case 2.  $k$  is odd.*

Then  $|X|$  is odd. If  $|X| = \gamma_{\text{pr}}(G) - 1$ , then  $|Y| \leq \gamma_{\text{pr}}(G) - k - 1 = |X - Z|$ .

Let  $\pi$  be any permutation of  $V$  such that

$$\pi(Y) \subseteq X - Z, \quad \pi(Z) = Z \quad \text{and} \quad Y \subseteq \pi(X - Z).$$

Then  $W = X_1 \cup X_2$  dominates  $\pi(G)$  and it is easy to see that  $\langle W \rangle$  has a perfect matching. Therefore  $W$  is a PDS of  $\pi G$  and

$$|W| = 2|X| = 2\gamma_{\text{pr}}(G) - 2,$$

so  $G$  is not a universal doubler.

Thus we assume that  $0 < |X| < \gamma_{\text{pr}}(G) - 2$ . Similar to Case 1, we choose the PDS  $D$  of  $G$  as follows.

- (iii) If  $|Y| + k \leq \gamma_{\text{pr}}(G)$ , let  $D$  be any  $\gamma_{\text{pr}}$ -set of  $G$ .
- (iv) Otherwise, let  $D$  be any PDS of  $G$  with  $|D| = |Y| + k - 1$  if  $|Y|$  is even, or  $|D| = |Y| + k$  if  $|Y|$  is odd.

Let  $w \in Z$  and let  $\pi$  be any permutation of  $V$  such that  $\pi(Y \cup Z - \{w\}) \subseteq D$ ,  $\pi(Z - \{w\})$  has a perfect matching  $M'$  which is contained in a  $D$ -matching, and  $\pi(w) = w' \in V - D$ . Let  $W = X_1 \cup D_2 \cup \{w'_2\}$ . Since  $X_1$  dominates  $G_1 - Y_1$  and  $D_2$  dominates  $G_2$  and  $Y_1$ , it follows that  $W$  dominates  $\pi G$ . Also,  $\langle W \rangle$  has a perfect matching in which  $w_1$  is paired with  $w'_2$ , and each edge  $u_2 v_2$  in  $M'_2$  is replaced by two edges  $z_1 u_2$  and  $z'_1 v_2$ , where  $z, z' \in Z - \{w\}$ . Therefore  $W$  is a PDS of  $G$ . If  $D$  was chosen in (iii) and thus is a  $\gamma_{\text{pr}}$ -set of  $G$ , then

$$|W| = |X| + |D| + 1 < \gamma_{\text{pr}}(G) - 2 + \gamma_{\text{pr}}(G) + 1 = 2\gamma_{\text{pr}}(G) - 1$$

and  $G$  is not a universal  $\gamma_{\text{pr}}$ -doubler. On the other hand, if  $D$  was chosen in (iv), then  $|D| \leq |Y| + k$ , so

$$\begin{aligned} |W| &= |X| + |D| + 1 \\ &< |X| + 2\gamma_{\text{pr}}(G) - |X| - k - 1 + k + 1 \\ &= 2\gamma_{\text{pr}}(G) \end{aligned}$$

and once again  $G$  is not a universal  $\gamma_{\text{pr}}$ -doubler.

Conversely, let  $\pi$  be a permutation of  $V$  such that  $\gamma_{\text{pr}}(\pi G) < 2\gamma_{\text{pr}}(G) - 1$  and consider any  $\gamma_{\text{pr}}$ -set  $W$  of  $\pi G$ . Define

$$X_1 = W \cap V_1 \quad \text{and} \quad D_2 = W \cap V_2.$$

Assume without loss of generality that  $|X_1| < \gamma_{\text{pr}}(G)$ . Let  $M'$  be a  $W$ -matching and let  $D'_2$  be the set of vertices in  $D_2$  which are not paired with another vertex in  $D_2$  under  $M'$ . Say  $|D'_2| = k'$ . Also, let  $k$  be the number of vertices not paired in a maximum matching of  $\langle X_1 \rangle$ . Note that  $k \leq k'$ .

If  $X_1 \neq \emptyset$ , then  $|D_2| < 2\gamma_{\text{pr}}(G) - |X| - 1$  and each vertex of  $D_2 - D'_2$  dominates at most one vertex in  $Y_1$ , while no vertex in  $D'_2$  dominates a vertex in  $Y_1$ . Therefore  $|Y_1| \leq |D_2 - D'_2|$ , which implies that

$$|Y| < 2\gamma_{\text{pr}}(G) - |X| - k' - 1 \leq 2\gamma_{\text{pr}}(G) - |X| - k - 1.$$

If  $X_1 = \emptyset$ , then  $D_2$  dominates  $V_1$  and so  $D_2 = V_2$ . Therefore  $n = |D_2| < 2\gamma_{\text{pr}}(G)$ , so that by Lemma 3,  $\gamma(G) < \gamma_{\text{pr}}(G)$ . Let  $X'$  be a  $\gamma$ -set of  $G$ ,  $Y' = V - N[X']$  and  $k'$  be the number of vertices not paired in a maximum matching of  $\langle X' \rangle$ . Since  $k' \leq |X'| < \gamma_{\text{pr}}(G)$ ,

$$(2) \quad |Y'| = 0 < 2\gamma_{\text{pr}}(G) - |X'| - k' - 1. \quad \blacksquare$$

As an example of universal  $\gamma_{\text{pr}}$ -doubblers, consider the following family  $\mathcal{F}$  of graphs. Form the graph  $F_{2n} \in \mathcal{F}$  by joining each vertex of  $C_{2n}$  to  $2n - 1$  new vertices. Note that  $\gamma_{\text{pr}}(F_{2n}) = \gamma(F_{2n}) = 2n$ . Figure 3 shows the graph  $F_4$ .




Figure 3.  $F_4 \in \mathcal{F}$ : An example of a universal  $\gamma_{\text{pr}}$ -doubler.

By Theorem 4, to prove that  $F_{2n}$  is a universal  $\gamma_{\text{pr}}$ -doubler, we must show that for each pair of sets  $X, Y \subseteq V(F_{2n})$  as defined in (1),  $|Y| \geq 2\gamma_{\text{pr}}(F_{2n}) - |X| - k - 1$ . Suppose  $|X| = 2n - d$ , where  $1 \leq d \leq 2n - 1$ . It is easy to see



that  $|Y| \geq d(2n - 1)$ . If  $d = 1$ , then  $k \geq 1$ , hence

$$2\gamma_{\text{pr}}(F_{2n}) - |X| - k - 1 \leq 4n - (2n - 1) - 1 - 1 = 2n - 1 \leq |Y|.$$

If  $2 \leq d \leq 2n - 1$ , then  $k \geq 0$ , hence

$$\begin{aligned} 2\gamma_{\text{pr}}(F_{2n}) - |X| - k - 1 &\leq 4n - (2n - d) - 1 \\ &= 2n + d - 1 \\ &\leq 2n + (2n - 1) - 1 \\ &= 2(2n - 1) \\ &\leq d(2n - 1) \\ &\leq |Y|. \end{aligned}$$

Note that to construct a universal  $\gamma_{\text{pr}}$ -doubler  $G$  from  $C_{2n}$  by adding pendant edges at vertices of  $C_{2n}$ , at least  $2n - 1$  pendant edges must be added at each vertex of  $C_{2n}$ . If some vertices of  $C_{2n}$  are joined to more than  $2n - 1$  new vertices, the resulting graph is also a universal  $\gamma_{\text{pr}}$ -doubler.

**Corollary 5.** *If  $\gamma(G) = \gamma_{\text{pr}}(G) = 2$ , then  $G$  is a universal  $\gamma_{\text{pr}}$ -doubler.*

**Proof.** Suppose  $\gamma(G) = \gamma_{\text{pr}}(G) = 2$ . Let  $x \in V$  and  $Y = V - N[x]$ . Since  $\gamma(G) = 2$ ,  $|Y| \geq 1$ . The result follows from Theorem 4. ■

We use Theorem 4 to obtain the promised results on the degrees and number of external private neighbours of the vertices in  $\gamma_{\text{pr}}$ -sets of a universal  $\gamma_{\text{pr}}$ -doubler.

**Corollary 6.** *Let  $G$  be a universal  $\gamma_{\text{pr}}$ -doubler and  $D$  any  $\gamma_{\text{pr}}$ -set of  $G$ . Then  $|\text{epn}(v, D)| \geq \gamma_{\text{pr}}(G) - 1$  for each  $v \in D$ .*

**Proof.** Let  $X = D - \{v\}$ . Then  $X \neq \emptyset$  because  $\gamma_{\text{pr}}(G) \geq 2$ , and  $k = 1$  because there is only one vertex in  $X$  that is not paired. By Theorem 4,

$$|V - N[X]| \geq 2\gamma_{\text{pr}}(G) - |X| - k - 1 = \gamma_{\text{pr}}(G) - 1.$$

Since  $D$  is a dominating set,  $v$  dominates  $V - N[X]$ . Moreover,  $v \notin V - N[X]$  because  $v$  is dominated by its partner in  $D$ . Hence  $\text{epn}(v, D) = V - N[X]$  and the result follows. ■

The converse of Corollary 6 is shown to be false by the counterexample in Figure 4. The black vertices form the set  $D$ , which is the only  $\gamma_{\text{pr}}$ -set of  $G$ , and for all  $v \in D$ ,  $|\text{epn}(v, D)| = 3 = \gamma_{\text{pr}}(G) - 1$ . Let  $X$  consist of the circled vertices. Then

$$|Y| = |V - N[X]| = 2 < 2\gamma_{\text{pr}}(G) - |X| - k - 1 = 3,$$

so by Theorem 4,  $G$  is not a universal  $\gamma_{\text{pr}}$ -doubler.




Figure 4. A counterexample to the converse of Corollary 6.

**Corollary 7.** *If  $G$  is a universal  $\gamma_{\text{pr}}$ -doubler and  $v \in V$  is contained in a  $\gamma_{\text{pr}}$ -set of  $G$ , then  $\deg v \geq \gamma_{\text{pr}}(G)$ .*

**Proof.** Suppose  $D$  is a  $\gamma_{\text{pr}}$ -set of  $G$  and  $v \in D$ . By Corollary 6,  $|\text{epn}(v, D)| \geq \gamma_{\text{pr}}(G) - 1$ . Since  $v$  is paired with some vertex in  $D$ , the result follows. ■

The complete graphs of order at least three show that the converse of Corollary 7 is not true.

**Corollary 8.** *If  $G$  is a universal  $\gamma_{\text{pr}}$ -doubler of order  $n$ , then  $\gamma_{\text{pr}}(G) \leq \sqrt{n}$ .*

**Proof.** By Corollary 7,  $\deg v \geq \gamma_{\text{pr}}(G)$  for any vertex  $v$  of any  $\gamma_{\text{pr}}$ -set  $D$  of  $G$ . Hence  $n \geq [\gamma_{\text{pr}}(G)]^2$ . ■

We conclude this section by obtaining a sufficient condition for regular graphs to be universal  $\gamma_{\text{pr}}$ -doublers. This allows us to construct a family of universal  $\gamma_{\text{pr}}$ -doublers.

The PDS  $D$  is an *efficient paired dominating set* (EPDS) if  $N(u) \cap N(v) = \emptyset$  for any two vertices  $u, v \in D$ .

**Lemma 9.** *If  $G$  is regular and has an EPDS  $D$ , then  $\gamma_{\text{pr}}(G) = |D|$ .*

**Proof.** Let  $X$  be a  $\gamma_{\text{pr}}$ -set of  $G$ . Then  $|X| \leq |D|$  and by regularity,  $n \leq r|X|$ . Since  $D$  is an EPDS,  $n = r|D|$ . Hence  $|D| \leq |X|$  and so  $D$  is a  $\gamma_{\text{pr}}$ -set of  $G$ . ■

**Corollary 10.** *If  $G$  is  $r$ -regular with  $r \geq \gamma_{\text{pr}}(G)$  and  $G$  has an EPDS, then  $G$  is a universal  $\gamma_{\text{pr}}$ -doubler.*

**Proof.** Let  $X \subseteq V$  with  $0 < |X| < \gamma_{\text{pr}}(G)$  and define  $Y$  and  $k$  as in (1). Then  $|N[X]| \leq r|X| + k$ . Since  $G$  has an EPDS,  $n = r\gamma_{\text{pr}}(G)$ . Then

$$|Y| \geq r\gamma_{\text{pr}}(G) - r|X| - k \geq \gamma_{\text{pr}}(G)(\gamma_{\text{pr}}(G) - |X|) - k.$$

If  $|X| = \gamma_{\text{pr}}(G) - 1$ , then

$$|Y| \geq \gamma_{\text{pr}}(G) - k = 2\gamma_{\text{pr}}(G) - |X| - k - 1,$$

and if  $|X| \leq \gamma_{\text{pr}}(G) - 2$ , then

$$|Y| \geq 2\gamma_{\text{pr}}(G) - k.$$

In either case the hypothesis of Theorem 4 is satisfied and it follows that  $G$  is a universal  $\gamma_{\text{pr}}$ -doubler. ■

Corollary 10 allows us to construct a family  $\mathcal{H}$  of regular universal  $\gamma_{\text{pr}}$ -doublers. Label the vertices of  $C_{2m}$  consecutively by  $u_1, v_1, u_2, v_2, \dots, u_m, v_m$ . Construct each  $H_{2m,r} \in \mathcal{H}$  by replacing alternate edges  $u_i v_i$ ,  $i = 1, \dots, m$ , of  $C_{2m}$  by a copy of  $B_i \cong K_{r-1, r-1}$ ,  $r \geq 2m$ , joining  $u_i$  to each vertex in one partite set, and  $v_i$  to each vertex in the other partite set of  $B_i$ . See Figure 5 for  $H_{4,4}$ .




Figure 5. The 4-regular universal  $\gamma_{\text{pr}}$ -doubler  $H_{4,4}$ .

Clearly,  $H_{2m,r}$  is  $r$ -regular. It is also easy to see that  $\bigcup_{i=1}^m \{u_i, v_i\}$  forms an efficient  $\gamma_{\text{pr}}$ -set of  $H_{2m,r}$  (in which each  $v_i$  is partnered by  $u_{i+1 \pmod m}$ ). By Corollary 10,  $H_{2m,r}$  is a universal  $\gamma_{\text{pr}}$ -doubler.

#### 4. PRISM DOUBLERS

It is reasonable to expect that there are graphs that are prism  $\gamma_{\text{pr}}$ -doublers but not universal  $\gamma_{\text{pr}}$ -doublers. In this section we first supply necessary and sufficient conditions in Theorem 11, and then a simpler sufficient condition in Proposition 12, for a graph to be a prism doubler. The latter result combined with Corollary 6 allows us to construct prism  $\gamma_{\text{pr}}$ -doublers that are not universal  $\gamma_{\text{pr}}$ -doublers.

**Theorem 11.** *A graph  $G$  is a prism  $\gamma_{\text{pr}}$ -doubler if and only if for each set  $X \subseteq V$  with  $0 < |X| < \gamma_{\text{pr}}(G)$ , any matching  $M$  of  $\langle X \rangle$ , and  $Y$  and  $k$  as defined in (1), either*

- (i)  $|Y| \geq 2\gamma_{\text{pr}}(G) - |X| - k - 1$ , or
- (ii)  $|Y| = 2\gamma_{\text{pr}}(G) - |X| - k - d - 1$ , where  $d \geq 1$ , and if  $A \subseteq N[X] - Z$  dominates  $N\{X\} - N[Y] - N[Z]$  and  $\langle A \cup Y \rangle$  has a perfect matching, then  $|A| \geq d$ .

**Proof.** Assume  $\gamma_{\text{pr}}(K_2 \square G) = 2\gamma_{\text{pr}}(G)$  and consider any pair of sets  $X, Y$  as defined in (1) and a matching  $M$  of  $\langle X \rangle$ . If  $|Y| \geq 2\gamma_{\text{pr}}(G) - |X| - k - 1$  then we are done, so assume  $|Y| = 2\gamma_{\text{pr}}(G) - |X| - k - d - 1$  for some  $d \geq 1$ .

Suppose to the contrary that there exists a set  $A \subseteq N[X] - Z$  such that  $A$  dominates  $N\{X\} - N[Y] - N[Z]$  and  $\langle A \cup Y \rangle$  has a perfect matching  $M^*$ , but  $|A| \leq d - 1$ . Define the set  $W \subseteq V(K_2 \square G)$  by  $W = X_1 \cup Y_2 \cup A_2 \cup Z_2$ . By the definition of  $X$  and  $Y$ ,  $X_1 \cup Y_2$  dominates  $G_1$ . Since  $A_2$  dominates  $N\{X_2\} - N[Y_2] - N[Z_2]$ ,  $W$  also dominates  $G_2$ . Thus  $W$  dominates  $K_2 \square G$ . Moreover,  $M \cup M^* \cup \{z_1 z_2 : z \in Z\}$  is a  $W$ -matching, so  $W$  is a PDS of  $K_2 \square G$ . But

$$\begin{aligned} |W| &= |X| + |Y| + |Z| + |A| \\ &\leq |X| + (2\gamma_{\text{pr}}(G) - |X| - k - d - 1) + k + (d - 1) \\ &= 2\gamma_{\text{pr}}(G) - 2, \end{aligned}$$

a contradiction. Thus (ii) holds.

Conversely, assume  $\gamma_{\text{pr}}(K_2 \square G) < 2\gamma_{\text{pr}}(G) - 1$  and let  $W = X_1 \cup D_2$  be a  $\gamma_{\text{pr}}$ -set of  $K_2 \square G$ . We may assume without loss of generality that  $|X| < \gamma_{\text{pr}}(G)$ .

We consider two cases, depending on whether  $X = \emptyset$  or  $X \neq \emptyset$ .

*Case 1.*  $X = \emptyset$ .

Then  $D_2 = V_2$  to dominate  $G_1$ . Therefore

$$|W| = |D| = n \leq 2\gamma_{\text{pr}}(G) - 2.$$

By Lemma 3,  $\gamma(G) < \gamma_{\text{pr}}(G)$ . Let  $X'$  be a  $\gamma$ -set of  $G$ ,  $M'$  be a maximum matching of  $\langle X' \rangle$ ,  $Z'$  the set of  $\overline{M}$ -vertices in  $X'$  and  $k' = |Z'|$ . Then  $k' > 0$  because  $X'$  is not a PDS of  $G$ , and  $Y' = V - N[X'] = \emptyset$  because  $X'$  dominates  $G$ . But

$$2\gamma_{\text{pr}}(G) - |X'| - k' - 1 \geq 2\gamma_{\text{pr}}(G) - 2|X'| - 1 > 0 = |Y'|$$

and so (i) does not hold. Hence there exists a positive integer  $d$  such that

$$0 = |Y'| = 2\gamma_{\text{pr}}(G) - |X'| - k' - d - 1,$$

$$\text{i.e., } d = 2\gamma_{\text{pr}}(G) - |X'| - k' - 1.$$

Let  $A' = X' - Z'$ . Then  $A' \subseteq N[X'] - Z'$ ,  $A'$  dominates  $N\{X'\} - N[Y'] - N[Z']$  and, since  $Y' = \emptyset$ ,  $M'$  is a perfect matching of  $\langle A' \cup Y' \rangle$ . But

$$|A'| = |X'| - k' = 2|X'| - |X'| - k' < 2\gamma_{\text{pr}}(G) - |X'| - k' - 1 = d,$$


thus (ii) also does not hold.

*Case 2.*  $X \neq \emptyset$ .

Let  $M^*$  be a  $W$ -matching, let  $M_1$  be the matching of  $\langle X_1 \rangle$  induced by  $M^*$ , and let  $Z_1$  be the set of vertices in  $X_1$  which are paired with vertices in  $D_2$  (i.e., the vertices in  $Z_2$ ) under  $M^*$ . Then in  $G$ ,  $Z$  is the set of  $\overline{M}$ -vertices in  $X$ , and  $Z \subseteq D$ . Define  $Y$  and  $k$  as in (1). Since  $D_2$  dominates  $Y_1, Y_2 \subseteq D_2$  and so  $Y \subseteq D$ . Moreover,  $Y \cap Z = \emptyset$ . Hence

$$|Y| \leq |D| - |Z| < 2\gamma_{\text{pr}}(G) - |X| - 1 - k.$$

Therefore (i) does not hold.

Figure 6.  $\langle A \cup Y \rangle$  has a perfect matching in  $G$ .

Let  $A = D - Z - Y$ . Then  $A \subseteq N[X] - Z$ . Since  $D_2$  dominates all vertices of  $G_2$  except possibly the vertices in  $X_2 - D_2$ ,  $D$  dominates  $N\{X\}$ , and so  $A$  dominates  $N\{X\} - N[Y] - N[Z]$ . Moreover,  $A \cup Y = D - Z$  and so  $\langle A \cup Y \rangle$  has a perfect matching (corresponding to the edges of  $M^*$  with both endvertices in  $D_2$ ). (See Figure 6, where the black vertices indicate  $X_1$  in  $G_1$ ,  $D_2$  in  $G_2$ , and  $A$  in  $G$ , the grey vertices indicate  $Y$  in  $G$ , and the dark edges indicate the matching  $M^*$  in  $K_2 \square G$  and the perfect matching in  $A \cup Y$ .) Since

$$Y = D - Z - A, \quad A \cup Z \subseteq D \quad \text{and} \quad A \cap Z = \emptyset,$$

it follows that

$$|Y| = |D| - |Z| - |A| < 2\gamma_{\text{pr}}(G) - 1 - |X| - k - |A|.$$

Thus

$$|Y| = 2\gamma_{\text{pr}}(G) - |X| - k - d - 1$$

for some  $d > |A|$ , and so (ii) also does not hold. ■

The following proposition enables us to describe classes of prism  $\gamma_{\text{pr}}$ -doubblers that are not universal  $\gamma_{\text{pr}}$ -doubblers.

**Proposition 12.** *If every vertex that is contained in a  $\gamma_{\text{pr}}$ -set of  $G \neq K_2$  is adjacent to at least one leaf, then  $G$  is a prism  $\gamma_{\text{pr}}$ -doubler.*

**Proof.** It is obvious that any support vertex of a graph  $G$  is contained in each PDS of  $G$ . Thus, if  $G$  satisfies the hypothesis, then  $\gamma_{\text{pr}}(G) = k$ , where  $k$  is the number of support vertices of  $G$ . Say  $u \in V$  is adjacent to the leaf  $v$ . Then in  $K_2 \square G$ ,  $u_1, v_1, v_2, u_2, u_1$  is an induced 4-cycle, and  $\deg v_1 = \deg v_2 = 2$ . Thus any PDS of  $K_2 \square G$  contains at least two of these vertices, so that  $\gamma_{\text{pr}}(K_2 \square G) \geq 2k$ , and the result follows. ■

Now let  $H$  be a graph of order  $k \geq 4$  that has a perfect matching and let  $G$  be any graph obtained by joining each vertex of  $H$  to at least one leaf, and some vertex  $v$  to at most  $k-2$  leaves. By Proposition 12,  $G$  is a prism  $\gamma_{\text{pr}}$ -doubler with  $\gamma_{\text{pr}}$ -set  $V(H)$  and  $\gamma_{\text{pr}}(G) = k$ . However, by Corollary 6,  $G$  is not a universal  $\gamma_{\text{pr}}$ -doubler, because  $|\text{epn}(v, V(H))| \leq k-2 < \gamma_{\text{pr}}(G) - 1 = k-1$ .

## 5. PRISM FIXERS

Since  $\gamma_{\text{pr}}(K_2 \square G) \leq 2\gamma(G)$  for any graph  $G$ , it is immediately clear that if  $\gamma_{\text{pr}}(G) = 2\gamma(G)$ , then  $G$  is a prism  $\gamma_{\text{pr}}$ -fixer. Examples of such graphs include nontrivial complete graphs,  $P_5$ ,  $C_5$  and  $C_6$ . We now extend this result to determine a necessary and sufficient condition for a graph to be a prism  $\gamma_{\text{pr}}$ -fixer.

Let  $S \subseteq V$  such that  $\langle S \rangle$  has a perfect matching  $M$ . A *paired partition* of  $S$  is a partition  $S_1, \dots, S_k$  such that each edge of  $M$  is contained in  $\langle S_i \rangle$  for some  $i$ . A *weak paired partition* is a paired partition in which some of the sets may be empty. A *split* of  $S$  is a partition  $S = S_1 \cup S_2$  such that each edge of  $M$  has one endvertex in  $S_1$  and the other one in  $S_2$ .

In our next theorem we consider a weak paired partition  $S = D \cup Y \cup Z$  of a  $\gamma_{\text{pr}}$ -set  $S$  of  $G$ , and define  $U = (V - S) \cap N[D] \cap N[Z]$  and  $X = V - S - U$ . Note that each vertex in  $U$  is adjacent to a vertex in  $D$  and to a vertex in  $Z$ , each vertex in  $X$  is adjacent to vertices in at most one of  $D$  and  $Z$ , and any vertex of  $G - S$  may or may not be adjacent to a vertex in  $Y$ . See Figure 7, where  $S$  consists of the black vertices,  $U$  of the grey vertices and  $X$  of the white vertices, and where the vertices in  $D$  are indicated by circles, those in  $Z$  by squares, and those in  $Y$  by triangles.

**Theorem 13.** *A graph  $G$  is a prism  $\gamma_{\text{pr}}$ -fixer if and only if  $G$  has a  $\gamma_{\text{pr}}$ -set  $S$  with a weak paired partition  $S = D \cup Y \cup Z$  in which  $Y$  has a split  $Y = Y' \cup Y''$  such that  $Y'$  dominates  $X = V - S - (N[D] \cap N[Z])$ .*




Figure 7. Examples of weak paired partitions.

**Proof.** Suppose  $G$  is a prism  $\gamma_{\text{pr}}$ -fixer and let  $W$  be a  $\gamma_{\text{pr}}$ -set of  $K_2 \square G$ . Say  $D'_1 = W \cap V_1$  and  $Z'_2 = W \cap V_2$ . Let  $M^*$  be a  $W$ -matching in which as few vertices as possible are matched with their own image. Let

$$S' = D' \cup Z',$$

$$Y' = D' \cap Z',$$

$M'$  be the matching of  $\langle S' \rangle$  induced by  $M^*$ ,

$R$  be the set of  $\overline{M'}$ -vertices.

Then  $S'$  dominates  $G$ ,  $R \subseteq Y'$ , and if  $u \in R$ , then  $u_1 u_2 \in E(M^*)$ . Say  $R = \{u^1, \dots, u^k\}$ , let  $S^0 = S'$  and for  $i = 1, \dots, k$ , construct  $S^i$  recursively as follows.

- (i) If  $u^i$  is adjacent to  $s^i \in V - S^{i-1}$ , let  $S^i = S^{i-1} \cup \{s^i\}$ .
- (ii) Otherwise,  $u^i$  is adjacent to some vertex in  $S^{i-1}$  because  $G$  is isolate-free, hence  $N[u^i] \subseteq S^{i-1}$ ; let  $S^i = S^{i-1} - \{u^i\}$ .

Then  $S^k$  dominates  $G$ ,  $\langle S^k \rangle$  has a perfect matching and thus  $S^k$  is a PDS of  $G$ . Moreover,

$$(3) \quad |S^k| \leq |D'| + |Z'| - |Y'| + |R| \leq |W|.$$

But  $G$  is a prism  $\gamma_{\text{pr}}$ -fixer, so equality holds in (3). In particular,  $R = Y'$  and each  $S^i$  is constructed as described in (i). Moreover,  $Y'$  is independent, for if  $u, v \in Y'$  and  $uv \in E$ , then  $u_1 u_2, v_1 v_2 \in E(M^*)$  (since  $u$  and  $v$  are  $\overline{M'}$ -vertices) and  $(M^* - \{u_1 u_2, v_1 v_2\}) \cup \{u_1 v_1, u_2 v_2\}$  is a  $W$ -matching in which



fewer vertices are mapped to their own images than in  $M^*$ , contradicting the choice of  $M^*$ .

Let

$$S = S^k, \quad D = D' - Y', \quad Z = Z - Y', \quad Y = S - D - Z \quad \text{and} \quad Y'' = Y - Y'.$$

Then  $D \cup Z \cup Y$  is a weak paired partition of  $S$  and  $Y' \cup Y''$  is a split of  $Y$  and we only need to prove that  $Y'$  dominates  $X$ . Suppose  $x \in X$ . We assume that  $x \notin N[D]$ ; the case  $x \notin N[Z]$  is similar. Since  $x \notin S$ ,  $x_1 \notin D'_1$  and  $x_2 \notin Z'_2$ . Thus  $x_1$  is dominated in  $G_1$  by a vertex in  $D'_1 - D_1$ , i.e., by a vertex in  $Y'_1$ . Therefore  $x$  is dominated by a vertex in  $Y'$  as required.

Conversely, assume  $G$  has a  $\gamma_{\text{pr}}$ -set  $S$  that satisfies the conditions of the theorem. Then  $D_1 \cup Y'_1$  dominates  $(G_1 - Z_1) \cup D_2$ , and  $Z_2 \cup Y'_2$  dominates  $(G_2 - D_2) \cup Z_1$ . Hence  $W = D_1 \cup Z_2 \cup Y'_1 \cup Y'_2$  is a PDS of  $K_2 \square G$  and  $|W| = |S| = \gamma_{\text{pr}}(G)$ . By Proposition 1,  $W$  is a  $\gamma_{\text{pr}}$ -set of  $K_2 \square G$ . ■

The three graphs in Figure 7 are examples of prism  $\gamma_{\text{pr}}$ -fixers. Other examples of prism fixers include  $K_n$  for  $n \geq 2$ ,  $P_n$  for  $n \in \{3, 5, 6, 9\}$  and  $C_n$  for  $n \in \{5, 6, 9\}$ . (This list contains all paths and cycles that are prism  $\gamma_{\text{pr}}$ -fixers.)

## 6. PROBLEMS

We conclude with open problems related to the above material. The graph  $G$  in Figure 1 illustrates that the paired domination number of a graph may exceed the paired domination number of some of its generalized prisms. Note that this graph is  $\gamma_{\text{pr}}$ -edge-critical, i.e.,  $\gamma_{\text{pr}}(G + e) < \gamma_{\text{pr}}(G)$  for each edge  $e \in E(\overline{G})$ . (See [5], for example.)

### Problem 1.

- (i) Characterize the class of graphs  $G$  with  $\gamma_{\text{pr}}(\pi G) < \gamma_{\text{pr}}(G)$  for some permutation  $\pi$  of  $V$ .
- (ii) If  $\gamma_{\text{pr}}(\pi G) < \gamma_{\text{pr}}(G)$  for some permutation  $\pi$  of  $V$ , what is  $\max_{\pi \in S_n} \{\gamma_{\text{pr}}(\pi G)\}$ ?
- (iii) What is  $\min_{\pi \in S_n} \{\gamma_{\text{pr}}(\pi G)/\gamma_{\text{pr}}(G)\}$ ?
- (iv) If  $\gamma_{\text{pr}}(\pi G) < \gamma_{\text{pr}}(G)$  for some permutation  $\pi$  of  $V$ , does it follow that  $G$  is  $\gamma_{\text{pr}}$ -edge-critical? (The converse is not true—consider  $C_5$ .)

For the usual domination number  $\gamma$ , it is still an open problem to find a nontrivial connected universal fixer, or to show that no such graph exists. The corresponding problem for the paired domination number (for graphs  $G$  with  $\gamma_{\text{pr}}(G) \leq \gamma_{\text{pr}}(\pi G)$  for all permutations  $\pi$  of  $V$ ) has not been studied at all. It is easy to see that nontrivial complete graphs are universal  $\gamma_{\text{pr}}$ -fixers, but none of the other graphs listed at the end of Section 5 is a universal  $\gamma_{\text{pr}}$ -fixer.

**Problem 2.** Prove or disprove Conjecture 1: The graphs  $\overline{K_n}$ ,  $n \geq 1$ , are the only universal  $\gamma$ -fixers.

**Problem 3.**

- (i) Characterize the class of universal  $\gamma_{\text{pr}}$ -fixers.
- (ii) Failing (i), find examples of noncomplete universal  $\gamma_{\text{pr}}$ -fixers.

#### REFERENCES

- [1] B. Brešar, M.A. Henning and D.F. Rall, *Paired-domination of Cartesian products of graphs*, Util. Math. **73** (2007) 255–265.
- [2] A.P. Burger and C.M. Mynhardt, *Regular graphs are not universal fixers*, Discrete Math. **310** (2010) 364–368.
- [3] A.P. Burger, C.M. Mynhardt and W.D. Weakley, *On the domination number of prisms of graphs*, Discuss. Math. Graph Theory **24** (2004) 303–318.
- [4] E.J. Cockayne, R.G. Gibson and C.M. Mynhardt, *Claw-free graphs are not universal fixers*, Discrete Math. **309** (2009) 128–133.
- [5] M. Edwards, R.G. Gibson, M.A. Henning and C.M. Mynhardt, *On paired-domination edge critical graphs*, Australasian J. Combin. **40** (2008) 279–292.
- [6] R.G. Gibson, *Bipartite graphs are not universal fixers*, Discrete Math. **308** (2008) 5937–5943.
- [7] B.L. Hartnell and D.F. Rall, *On Vizing’s conjecture*, Congr. Numer. **82** (1991) 87–96.
- [8] B.L. Hartnell and D.F. Rall, *On dominating the Cartesian product of a graph and  $K_2$* , Discuss. Math. Graph Theory **24** (2004) 389–402.
- [9] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker, New York, 1998).

- [10] C.M. Mynhardt and Z. Xu, *Domination in prisms of graphs: Universal fixers*, Utilitas Math. **78** (2009) 185–201.
- [11] M. Schurch, Domination Parameters for Prisms of Graphs (Master’s thesis, University of Victoria, 2005).

Received 29 January 2009

Revised 27 July 2009

Accepted 27 July 2009