

PROBLÈMES

P 844, R 1. In case $v = 2$ the solution was given by W. Narkiewicz and F. Rayner⁽¹⁾. An algorithm leading to a solution of the problem for every particular value of $v \geq 3$ was given by W. Narkiewicz⁽²⁾. The case $v = 3$ was there treated in detail. This algorithm was implemented by F. Rayner⁽³⁾ who determined the integers in question for all $v \leq 107$.

XXVII.2, p. 290

⁽¹⁾ W. Narkiewicz and F. Rayner, *Distribution of values of $\sigma_2(n)$ in residue classes*, Monatshefte für Mathematik 94 (1982), p. 133–141.

⁽²⁾ W. Narkiewicz, *Distribution of coefficients of Eisenstein series in residue classes*, Acta Arithmetica 43 (1983), p. 83–92.

⁽³⁾ F. Rayner, *Weak uniform distribution for divisor functions*, to appear.

P 1260, R 1. Professor V. Klee and Dr. P. Krupski have kindly informed us that the theorem, to which the problem pertains⁽⁴⁾, has been known since long. It has been stated by K. Menger⁽⁵⁾ and proved by F. Alt and G. Beer⁽⁶⁾. Another proof follows from a result of I. J. Schoenberg⁽⁷⁾. V. Klee has generalized the theorem and answered P 1260 in the negative⁽⁸⁾.

XLVI. 2, p. 204

Letter of February 18, 1983

⁽⁴⁾ J. Väisälä, *Dividing an arc to subarcs with equal chords*, this journal 46 (1982), p. 203–204.

⁽⁵⁾ K. Menger, *Mathematische Annalen* 103 (1930), p. 466–501.

⁽⁶⁾ F. Alt and G. Beer, *Ergebnisse eines mathematischen Kolloquiums* 6 (1935), p. 7.

⁽⁷⁾ I. J. Schoenberg, *Annals of Mathematics* 41 (1940), p. 715–720.

⁽⁸⁾ V. Klee, *Ratio-sequences of chains in connected metric spaces in: The geometry of metric and linear spaces*, Proceedings of the Conference, Michigan State University 1974, Lecture Notes in Mathematics 490, Springer-Verlag, Berlin 1975, p. 134–146.

THEMISTOCLES M. RASSIAS (ATHENS)

P 1284. Is the following true: there is no harmonic homeomorphism of the open unit ball B in R^3 , i.e. there are no harmonic functions f_1, f_2, f_3 defined in $B = \{\zeta = (\zeta_1, \zeta_2, \zeta_3) : |\zeta| < 1\}$, such that the mapping $\zeta \rightarrow (f_1, f_2, f_3)$ is a homeomorphism of B onto all of R^3 ?

MARTIN GA VALEC (KOŠICE)

P 1285. Formulé dans la commutation *Iterated products of ideals of Borel sets.*

Ce fascicule, p. 51

JAN VAN MILL AND EVERT WATTEL (AMSTERDAM)

P 1286. Formulé dans la communication *Partitioning spaces in homeomorphic rigid parts.*

Ce fascicule, p. 101
